

# COMMON ENTRANCE TEST - 2010

DATE	SUBJECT	TIME
<b>28-04-2010</b>	<b>MATHEMATICS</b>	<b>02.30 PM to 03.50 PM</b>

MAXIMUM MARKS	TOTAL DURATION	MAXIMUM TIME FOR ANSWERING
<b>60</b>	<b>80 MINUTES</b>	<b>70 MINUTES</b>

MENTION YOUR CET NUMBER	QUESTION BOOKLET DETAILS	
	VERSION CODE	SERIAL NUMBER
	<b>A - 1</b>	<b>378753</b>

**DOs :**

1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
2. This Question Booklet is issued to you by the Invigilator after the 2<sup>nd</sup> Bell, i.e., after **02.30 p.m.**
3. The Serial Number of this question booklet should be entered on the OMR answer sheet.
4. The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
5. Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

**DON'Ts :**

1. **THE TIMING AND MARKS PRINTED ON THE OMR ANSWER SHEET SHOULD NOT BE DAMAGED/ MUTILATED / SPOILED.**
2. Until the 3<sup>rd</sup> Bell is rung at **02.40 p.m.** :
  - Do not remove the seal/staple present on the right hand side of this question booklet.
  - Do not look inside this question booklet.
  - Do not start answering on the OMR answer sheet.

### IMPORTANT INSTRUCTIONS TO CANDIDATES

1. This question booklet contains 60 questions and each question will have four different options / choices.
2. After the 3<sup>rd</sup> Bell is rung at **02.40 p.m.**, remove the seal/staple present on the right hand side of this question booklet and start answering on the OMR answer sheet.
3. During the subsequent 70 minutes :
  - Read each question carefully.
  - Choose the correct answer from out of the four available options / choices given under each question.
  - **Completely darken/shade the relevant circle with a BLUE OR BLACK INK BALLPOINT PEN against the question number on the OMR answer sheet.**

**CORRECT METHOD OF SHADING THE CIRCLE ON THE OMR SHEET IS AS SHOWN BELOW :**



4. Please note that even a minute unintended ink dot on the OMR sheet will also be recognized and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
5. Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
6. After the **last bell** is rung at **03.50 p.m.**, stop writing on the OMR answer sheet and affix your **LEFT HAND THUMB IMPRESSION** on the OMR answer sheet as per the instructions.
7. Hand over the OMR ANSWER SHEET to the room Invigilator as it is.
8. After separating and retaining the top sheet (KEA Copy), the Invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.



6. If  $a, -a, b$  are the roots of  $x^3 - 5x^2 - x + 5 = 0$ , then  $b$  is a root of .....
- |                        |                        |
|------------------------|------------------------|
| 1) $x^2 - 3x - 10 = 0$ | 2) $x^2 + 5x - 30 = 0$ |
| 3) $x^2 + 3x - 20 = 0$ | 4) $x^2 - 5x + 10 = 0$ |
7. In the binomial expansion of  $(1+x)^{15}$ , the coefficients of  $x^r$  and  $x^{r+3}$  are equal. Then  $r$  is .....
- |      |      |
|------|------|
| 1) 4 | 2) 6 |
| 3) 8 | 4) 7 |
8. The  $n^{\text{th}}$  term of the series  $1 + 3 + 7 + 13 + 21 + \dots$  is 9901. The value of  $n$  is .....
- |        |       |
|--------|-------|
| 1) 900 | 2) 99 |
| 3) 100 | 4) 90 |
9. If  $\frac{1}{(3-5x)(2+3x)} = \frac{A}{3-5x} + \frac{B}{2+3x}$ , then  $A : B$  is .....
- |          |          |
|----------|----------|
| 1) 3 : 5 | 2) 3 : 2 |
| 3) 2 : 3 | 4) 5 : 3 |
10. Which of the following is NOT true?
- 1)  $p \rightarrow (q \wedge r) \equiv (p \rightarrow q) \wedge (p \rightarrow r)$ .
  - 2)  $\sim(p \leftrightarrow q) \equiv (p \wedge \sim q) \vee (\sim p \wedge q)$ .
  - 3)  $(p \wedge \sim q) \leftrightarrow (p \rightarrow q)$  is a tautology.
  - 4)  $\{(p \rightarrow q) \wedge (q \rightarrow r)\} \rightarrow (p \rightarrow r)$  is a tautology.

---

(Space for Rough Work)





21. If  $a > b > 0$ ,  $\text{Sec}^{-1}\left(\frac{a+b}{a-b}\right) = 2\text{Sin}^{-1}x$ , then  $x = \dots\dots\dots$

1)  $-\sqrt{\frac{a}{a+b}}$

2)  $\sqrt{\frac{a}{a+b}}$

3)  $-\sqrt{\frac{b}{a+b}}$

4)  $\sqrt{\frac{b}{a+b}}$

22. If  $x \neq n\pi$ ,  $x \neq (2n+1)\frac{\pi}{2}$ ,  $n \in Z$ , then  $\frac{\text{Sin}^{-1}(\text{Cos } x) + \text{Cos}^{-1}(\text{Sin } x)}{\text{Tan}^{-1}(\text{Cot } x) + \text{Cot}^{-1}(\text{Tan } x)} = \dots\dots\dots$

1)  $\frac{\pi}{4}$

2)  $\frac{\pi}{3}$

3)  $\frac{\pi}{2}$

4)  $\frac{\pi}{6}$

23. The general solution of  $1 + \text{Sin}^2 x = 3\text{Sin } x \cdot \text{Cos } x$ ,  $\text{Tan } x \neq \frac{1}{2}$  is  $\dots\dots\dots$

1)  $n\pi - \frac{\pi}{4}$ ,  $n \in Z$

2)  $n\pi + \frac{\pi}{4}$ ,  $n \in Z$

3)  $2n\pi + \frac{\pi}{4}$ ,  $n \in Z$

4)  $2n\pi - \frac{\pi}{4}$ ,  $n \in Z$

24. The least positive integer  $n$ , for which  $\frac{(1+i)^n}{(1-i)^{n-2}}$  is positive, is  $\dots\dots\dots$

1) 1

2) 2

3) 3

4) 4

25. If  $x + iy = (-1 + i\sqrt{3})^{2010}$ , then  $x = \dots\dots\dots$

1) 1

2) -1

3)  $-2^{2010}$

4)  $2^{2010}$

---

(Space for Rough Work)









41. If  $x = a \cos^3 \theta$  and  $y = a \sin^3 \theta$ , then  $\frac{dy}{dx} = \dots\dots\dots$

1)  $-\sqrt[3]{\frac{x}{y}}$

2)  $-\sqrt[3]{\frac{y}{x}}$

3)  $\sqrt[3]{\frac{y}{x}}$

4)  $\sqrt[3]{\frac{x}{y}}$

42. If  $y = \tan^{-1} \sqrt{x^2 - 1}$ , then the ratio  $\frac{d^2y}{dx^2} : \frac{dy}{dx} = \dots\dots\dots$

1)  $\frac{1+2x^2}{x(x^2+1)}$

2)  $\frac{x(x^2+1)}{1-2x^2}$

3)  $\frac{x(x^2-1)}{1+2x^2}$

4)  $\frac{1-2x^2}{x(x^2-1)}$

43.  $P$  is the point of contact of the tangent from the origin to the curve  $y = \log_e x$ . The length of the perpendicular drawn from the origin to the normal at  $P$  is .....

1)  $2\sqrt{e^2+1}$

2)  $\sqrt{e^2+1}$

3)  $\frac{1}{2e}$

4)  $\frac{1}{e}$

44. For the curve  $4x^5 = 5y^4$ , the ratio of the cube of the subtangent at a point on the curve to the square of the subnormal at the same point is .....

1)  $\left(\frac{4}{5}\right)^4$

2)  $\left(\frac{5}{4}\right)^4$

3)  $x\left(\frac{4}{5}\right)^4$

4)  $y\left(\frac{5}{4}\right)^4$

45. The set of real values of  $x$  for which  $f(x) = \frac{x}{\log x}$  is increasing, is .....

1)  $\{x : x < e\}$

2)  $\{1\}$

3)  $\{x : x \geq e\}$

4) empty

---

(Space for Rough Work)



51. The condition for the line  $y = mx + c$  to be a normal to the parabola  $y^2 = 4ax$  is .....

1)  $c = \frac{a}{m}$

2)  $c = 2am + am^3$

3)  $c = -2am - am^3$

4)  $c = -\frac{a}{m}$

52. The eccentric angle of the point  $(2, \sqrt{3})$  lying on  $\frac{x^2}{16} + \frac{y^2}{4} = 1$  is .....

1)  $\frac{\pi}{3}$

2)  $\frac{\pi}{6}$

3)  $\frac{\pi}{4}$

4)  $\frac{\pi}{2}$

53. The distance of the focus of  $x^2 - y^2 = 4$ , from the directrix which is nearer to it, is .....

1)  $2\sqrt{2}$

2)  $\sqrt{2}$

3)  $4\sqrt{2}$

4)  $8\sqrt{2}$

54. If  $\int f(x) \sin x \cdot \cos x \, dx = \frac{1}{2(b^2 - a^2)} \log f(x) + c$ , where  $c$  is the constant of integration,

then  $f(x) = \dots\dots\dots$

1)  $\frac{2}{ab \sin 2x}$

2)  $\frac{2}{(b^2 - a^2) \sin 2x}$

3)  $\frac{2}{ab \cos 2x}$

4)  $\frac{2}{(b^2 - a^2) \cos 2x}$

55. If  $\int \frac{\sqrt{x}}{x(x+1)} \, dx = k \tan^{-1} m$ , then  $(k, m)$  is .....

1)  $(1, \sqrt{x})$

2)  $(2, \sqrt{x})$

3)  $(2, x)$

4)  $(1, x)$

---

(Space for Rough Work)

56.  $\int_0^{\pi/4} \frac{\sin x + \cos x}{3 + \sin 2x} dx = \dots\dots\dots$

1)  $\frac{1}{2} \text{Log } 3$

2)  $2 \text{Log } 3$

3)  $\frac{1}{4} \text{Log } 3$

4)  $\text{Log } 3$

57.  $\int_0^1 x(1-x)^{3/2} dx = \dots\dots\dots$

1)  $\frac{24}{35}$

2)  $\frac{-8}{35}$

3)  $\frac{-2}{35}$

4)  $\frac{4}{35}$

58. The area bounded by the curve  $y = \begin{cases} x^2, & x < 0 \\ x, & x \geq 0 \end{cases}$  and the line  $y = 4$  is  $\dots\dots\dots$

1)  $\frac{40}{3}$

2)  $\frac{16}{3}$

3)  $\frac{32}{3}$

4)  $\frac{8}{3}$

59. The order and degree of the differential equation  $y = \frac{dp}{dx}x + \sqrt{a^2p^2 + b^2}$  where  $p = \frac{dy}{dx}$  (here  $a$  and  $b$  are arbitrary constants) respectively are  $\dots\dots\dots$

1) 1, 2

2) 2, 1

3) 2, 2

4) 1, 1

60. The general solution of the differential equation  $2x \frac{dy}{dx} - y = 3$  is a family of  $\dots\dots\dots$

1) straight lines

2) circles

3) hyperbolas

4) parabolas

---

(Space for Rough Work)