

<b>SUBJECT : PHYSICS</b>	<b>DAY-2</b>
<b>SESSION : MORNING</b>	<b>TIME : 10.30 A.M. TO 11.50 A.M.</b>

<b>MAXIMUM MARKS</b>	<b>TOTAL DURATION</b>	<b>MAXIMUM TIME FOR ANSWERING</b>
<b>60</b>	<b>80 MINUTES</b>	<b>70 MINUTES</b>

<b>MENTION YOUR CET NUMBER</b>	<b>QUESTION BOOKLET DETAILS</b>	
	<b>VERSION CODE</b>	<b>SERIAL NUMBER</b>
	<b>A - 1</b>	<b>470465</b>

**DOs :**

1. Check whether the CET No. has been entered and shaded in the respective circles on the OMR answer sheet.
2. This Question Booklet is issued to you by the invigilator after the **2<sup>nd</sup> Bell i.e., after 10.30 a.m.**
3. The Serial Number of this question booklet should be entered on the OMR answer sheet.
4. The Version Code of this question booklet should be entered on the OMR answer sheet and the respective circles should also be shaded completely.
5. Compulsorily sign at the bottom portion of the OMR answer sheet in the space provided.

**DON'TS :**

1. **THE TIMING AND MARKS PRINTED ON THE OMR ANSWER SHEET SHOULD NOT BE DAMAGED / MUTILATED / SPOILED.**
2. The **3<sup>rd</sup> Bell rings at 10.40 a.m., till then:**
  - Do not remove the paper seal present on the right hand side of this question booklet.
  - Do not look inside this question booklet.
  - Do not start answering on the OMR answer sheet.

**IMPORTANT INSTRUCTIONS TO CANDIDATES**

1. This question booklet contains 60 questions and each question will have one statement and four distracters. (Four different options / choices.)
2. After the **3<sup>rd</sup> Bell is rung at 10.40 a.m.,** remove the paper seal on the right hand side of this question booklet and check that this booklet does not have any unprinted or torn or missing pages or items etc., if so, get it replaced by a complete test booklet. Read each item and start answering on the OMR answer sheet.
3. During the subsequent 70 minutes:
  - Read each question carefully.
  - Choose the correct answer from out of the four available distracters (options / choices) given under each question / statement.
  - **Completely darken / shade the relevant circle with a BLUE OR BLACK INK BALL POINT PEN against the question number on the OMR answer sheet.**

**Correct Method of shading the circle on the OMR answer sheet is as shown below :**



4. Please note that even a minute unintended ink dot on the OMR answer sheet will also be recognised and recorded by the scanner. Therefore, avoid multiple markings of any kind on the OMR answer sheet.
5. Use the space provided on each page of the question booklet for Rough Work. Do not use the OMR answer sheet for the same.
6. After the **last bell is rung at 11.50 a.m.,** stop writing on the OMR answer sheet and affix your **LEFT HAND THUMB IMPRESSION** on the OMR answer sheet as per the instructions.
7. Hand over the **OMR ANSWER SHEET** to the room invigilator as it is.
8. After separating the top sheet (Our Copy), the invigilator will return the bottom sheet replica (Candidate's copy) to you to carry home for self-evaluation.
9. Preserve the replica of the OMR answer sheet for a minimum period of ONE year.

**P**



**[Turn Over**



4. A stone of mass 0.05 kg is thrown vertically upwards. What is the direction and magnitude of net force on the stone during its upward motion ?
- (1) 0.49 N vertically upwards
  - (2) 0.49 N vertically downwards
  - (3) 0.98 N vertically downwards
  - (4) 9.8 N vertically downwards
5. The kinetic energy of a body of mass 4 kg and momentum 6 Ns will be
- (1) 2.5 J
  - (2) 3.5 J
  - (3) 4.5 J
  - (4) 5.5 J
6. The ratio of angular speed of a second-hand to the hour-hand of a watch is
- (1) 720 : 1
  - (2) 60 : 1
  - (3) 3600 : 1
  - (4) 72 : 1
7. If the mass of a body is M on the surface of the earth, the mass of the same body on the surface of the moon is
- (1) M/6
  - (2) M
  - (3) 6 M
  - (4) Zero
8. Moment of Inertia of a thin uniform rod rotating about the perpendicular axis passing through its centre is I. If the same rod is bent into a ring and its moment of inertia about its diameter is I', then the ratio  $\frac{I}{I'}$  is
- (1)  $3/2 \pi^2$
  - (2)  $8/3 \pi^2$
  - (3)  $2/3 \pi^2$
  - (4)  $5/3 \pi^2$

---

**Space For Rough Work**






19. Three point charges  $3\text{nC}$ ,  $6\text{nC}$  and  $9\text{nC}$  are placed at the corners of an equilateral triangle of side  $0.1\text{ m}$ . The potential energy of the system is
- (1)  $8910\text{ J}$  (2)  $89100\text{ J}$   
(3)  $9910\text{ J}$  (4)  $99100\text{ J}$
20. A spherical shell of radius  $10\text{ cm}$  is carrying a charge  $q$ . If the electric potential at distances  $5\text{ cm}$ ,  $10\text{ cm}$  and  $15\text{ cm}$  from the centre of the spherical shell is  $V_1$ ,  $V_2$  and  $V_3$  respectively, then
- (1)  $V_1 > V_2 > V_3$  (2)  $V_1 < V_2 < V_3$   
(3)  $V_1 = V_2 > V_3$  (4)  $V_1 = V_2 < V_3$
21. A parallel plate capacitor is charged and then isolated. The effect of increasing the plate separation on charge, potential and capacitance respectively are
- (1) constant, decreases, decreases  
(2) increases, decreases, decreases  
(3) constant, decreases, increases  
(4) constant, increases, decreases
22. Four identical cells of emf  $E$  and internal resistance  $r$  are to be connected in series. Suppose if one of the cell is connected wrongly, the equivalent emf and effective internal resistance of the combination is
- (1)  $4E$  and  $4r$  (2)  $4E$  and  $2r$   
(3)  $2E$  and  $4r$  (4)  $2E$  and  $2r$
23. Three resistances  $2\Omega$ ,  $3\Omega$  and  $4\Omega$  are connected in parallel. The ratio of currents passing through them when a potential difference is applied across its ends will be
- (1)  $6 : 3 : 2$  (2)  $6 : 4 : 3$   
(3)  $5 : 4 : 3$  (4)  $4 : 3 : 2$

---


Space For Rough Work

24. Two cells of emf  $E_1$  and  $E_2$  are joined in opposition (such that  $E_1 > E_2$ ). If  $r_1$  and  $r_2$  be the internal resistance and  $R$  be the external resistance, then the terminal potential difference is



- (1)  $\frac{E_1 + E_2}{r_1 + r_2} \times R$                       (2)  $\frac{E_1 + E_2}{r_1 + r_2 + R} \times R$
- (3)  $\frac{E_1 - E_2}{r_1 + r_2} \times R$                       (4)  $\frac{E_1 - E_2}{r_1 + r_2 + R} \times R$

25. In the circuit shown below, the ammeter and the voltmeter readings are 3 A and 6 V respectively. Then the value of the resistance  $R$  is



- (1)  $2 \Omega$     (2)  $> 2 \Omega$
- (3)  $< 2 \Omega$     (4)  $\geq 2 \Omega$

---

Space For Rough Work

26. In Wheatstones network  $P = 2 \Omega$ ,  $Q = 2 \Omega$ ,  $R = 2 \Omega$  and  $S = 3 \Omega$ . The resistance with which S is to shunted in order that the bridge may be balanced is
- (1)  $1 \Omega$  (2)  $2 \Omega$   
(3)  $4 \Omega$  (4)  $6 \Omega$
27. The resistance of the bulb filament is  $100 \Omega$  at a temperature of  $100^\circ\text{C}$ . If its temperature co-efficient of resistance be  $0.005$  per  $^\circ\text{C}$ , its resistance will become  $200 \Omega$  at a temperature
- (1)  $300^\circ\text{C}$  (2)  $400^\circ\text{C}$   
(3)  $500^\circ\text{C}$  (4)  $200^\circ\text{C}$
28. Two concentric coils each of radius equal to  $2\pi$  cm are placed right angles to each other. If  $3\text{A}$  and  $4\text{A}$  are the currents flowing through the two coils respectively. The magnetic induction (in  $\text{Wb m}^{-2}$ ) at the centre of the coils will be
- (1)  $12 \times 10^{-5}$  (2)  $10^{-5}$   
(3)  $5 \times 10^{-5}$  (4)  $7 \times 10^{-5}$
29. A proton beam enters a magnetic field of  $10^{-4} \text{Wb m}^{-2}$  normally. If the specific charge of the proton is  $10^{11} \text{C kg}^{-1}$  and its velocity is  $10^9 \text{ms}^{-1}$ , then the radius of the circle described will be
- (1)  $0.1 \text{m}$  (2)  $10 \text{m}$   
(3)  $100 \text{m}$  (4)  $1 \text{m}$

---

Space For Rough Work



30. A cyclotron is used to accelerate
- (1) neutron
  - (2) only positively charged particles
  - (3) only negatively charged particles
  - (4) both positively and negatively charged particles
31. A galvanometer of resistance  $50 \Omega$  gives a full scale deflection for a current  $5 \times 10^{-4}$  A. The resistance that should be connected in series with the galvanometer to read 3 V is
- (1)  $595 \Omega$
  - (2)  $5050 \Omega$
  - (3)  $5059 \Omega$
  - (4)  $5950 \Omega$
32. Two parallel wires 1 m apart carry currents of 1 A and 3 A respectively in opposite directions. The force per unit length acting between these two wires is
- (1)  $6 \times 10^{-7} \text{ Nm}^{-1}$  repulsive
  - (2)  $6 \times 10^{-7} \text{ Nm}^{-1}$  attractive
  - (3)  $6 \times 10^{-5} \text{ Nm}^{-1}$  repulsive
  - (4)  $6 \times 10^{-5} \text{ Nm}^{-1}$  attractive
33. If there is no torsion in the suspension thread, then the time period of a magnet executing SHM is
- (1)  $T = \frac{1}{2\pi} \sqrt{\frac{MB}{I}}$
  - (2)  $T = \frac{1}{2\pi} \sqrt{\frac{I}{MB}}$
  - (3)  $T = 2\pi \sqrt{\frac{I}{MB}}$
  - (4)  $T = 2\pi \sqrt{\frac{MB}{I}}$
34. Core of electromagnets are made of ferromagnetic material which has
- (1) high permeability and low retentivity
  - (2) high permeability and high retentivity
  - (3) low permeability and high retentivity
  - (4) low permeability and low retentivity

---

Space For Rough Work

35. The magnetic susceptibility of a paramagnetic material at  $-73\text{ }^{\circ}\text{C}$  is 0.0075 and its value at  $-173\text{ }^{\circ}\text{C}$  will be
- (1) 0.0045 (2) 0.0030  
(3) 0.015 (4) 0.0075
36. Two coils have a mutual inductance 0.005 H. The current changes in the first coil according to the equation  $i = i_m \sin \omega t$  where  $i_m = 10\text{ A}$  and  $\omega = 100\pi\text{ rad s}^{-1}$ . The maximum value of the emf induced in the second coil is
- (1)  $2\pi$  (2)  $5\pi$   
(3)  $\pi$  (4)  $4\pi$
37. An aircraft with a wingspan of 40 m flies with a speed of 1080 km/hr in the eastward direction at a constant altitude in the northern hemisphere, where the vertical component of the earth's magnetic field  $1.75 \times 10^{-5}\text{ T}$ . Then the emf developed between the tips of the wings is
- (1) 0.5 V (2) 0.34 V  
(3) 0.21 V (4) 2.1 V
38. In an LCR circuit, at resonance
- (1) the current and voltage are in phase  
(2) the impedance is maximum  
(3) the current is minimum  
(4) the current leads the voltage by  $\pi/2$
39. A transformer is used to light 100 W – 110 V lamp from 220 V mains. If the main current is 0.5 A, the efficiency of the transformer is
- (1) 90% (2) 95%  
(3) 96% (4) 99%

---

Space For Rough Work

40. The average power dissipated in a pure inductor is

- (1)  $\frac{1}{2} VI$  (2)  $VI^2$   
(3)  $\frac{VI^2}{4}$  (4) zero

41. If  $\epsilon_0$  and  $\mu_0$  are the permittivity and permeability of free space and  $\epsilon$  and  $\mu$  are the corresponding quantities for a medium, then refractive index of the medium is

- (1)  $\sqrt{\frac{\mu_0 \epsilon_0}{\mu \epsilon}}$  (2)  $\sqrt{\frac{\mu \epsilon}{\mu_0 \epsilon_0}}$   
(3) 1 (4) Insufficient information

42. A person wants a real image of his own, 3 times enlarged. Where should he stand in front of a concave mirror of radius of curvature 30 cm ?

- (1) 10 cm (2) 30 cm  
(3) 90 cm (4) 20 cm

43. Calculate the focal length of a reading glass of a person if his distance of distinct vision is 75 cm.

- (1) 25.6 cm (2) 37.5 cm  
(3) 75.2 cm (4) 100.4 cm

44. In a Young's double slit experiment the slit separation is 0.5 m from the slits. For a monochromatic light of wavelength 500 nm, the distance of 3<sup>rd</sup> maxima from 2<sup>nd</sup> minima on the other side is

- (1) 2.75 mm (2) 2.5 mm  
(3) 22.5 mm (4) 2.25 mm

---

Space For Rough Work

45. To observe diffraction, the size of the obstacle
- (1) has no relation to wavelength.
  - (2) should be  $\lambda/2$ , where  $\lambda$  is the wavelength.
  - (3) should be much larger than the wavelength.
  - (4) should be of the order of wavelength.
46. The polarizing angle of glass is  $57^\circ$ . A ray of light which is incident at this angle will have an angle of refraction as
- (1)  $25^\circ$
  - (2)  $33^\circ$
  - (3)  $43^\circ$
  - (4)  $38^\circ$
47. Light of two different frequencies whose photons have energies 1 eV and 2.5 eV respectively, successively illuminate a metallic surface whose work function is 0.5 eV. Ratio of maximum speeds of emitted electrons will be
- (1) 1 : 5
  - (2) 1 : 4
  - (3) 1 : 2
  - (4) 1 : 1
48. Find the de-Broglie wavelength of an electron with kinetic energy of 120 eV.
- (1) 95 pm
  - (2) 102 pm
  - (3) 112 pm
  - (4) 124 pm
49. An  $\alpha$ -particle of energy 5 MeV is scattered through  $180^\circ$  by gold nucleus. The distance of closest approach is of the order of
- (1)  $10^{-10}$  cm
  - (2)  $10^{-12}$  cm
  - (3)  $10^{-14}$  cm
  - (4)  $10^{-16}$  cm

---

**Space For Rough Work**

50. If an electron in hydrogen atom jumps from an orbit of level  $n = 3$  to an orbit of level  $n = 2$ , the emitted radiation has a frequency ( $R = \text{Rydberg constant}$ ,  $C = \text{velocity of light}$ )

(1)  $\frac{3RC}{27}$

(2)  $\frac{RC}{25}$

(3)  $\frac{8RC}{9}$

(4)  $\frac{5RC}{36}$

51. What is the wavelength of light for the least energetic photon emitted in the Lyman series of the hydrogen spectrum. (take  $hc = 1240 \text{ eV nm}$ )

(1) 82 nm

(2) 102 nm

(3) 122 nm

(4) 150 nm

52. A nucleus at rest splits into two nuclear parts having radii in the ratio 1 : 2. Their velocities are in the ratio

(1) 8 : 1

(2) 6 : 1

(3) 4 : 1

(4) 2 : 1

53. The half life of a radioactive substance is 20 minutes. The time taken between 50 % decay and 87.5 % decay of the substance will be

(1) 30 minutes

(2) 40 minutes

(3) 25 minutes

(4) 10 minutes

54. A radioactive decay can form an isotope of the original nucleus with the emission of particles

(1) one  $\alpha$  and four  $\beta$

(2) one  $\alpha$  and two  $\beta$


(3) one  $\alpha$  and one  $\beta$

(4) four  $\alpha$  and one  $\beta$

---

Space For Rough Work

55. An LED is constructed from a pn junction based on a certain semi-conducting material whose energy gap is 1.9 eV. Then the wavelength of the emitted light is
- (1)  $2.9 \times 10^{-9}$  m                      (2)  $1.6 \times 10^{-8}$  m  
 (3)  $6.5 \times 10^{-7}$  m                      (4)  $9.1 \times 10^{-5}$  m
56. Amplitude modulation has
- (1) one carrier with two side band frequencies  
 (2) one carrier  
 (3) one carrier with infinite frequencies  
 (4) one carrier with high frequency
57. The circuit has two oppositely connected ideal diodes in parallel. What is the current flowing in the circuit ?



- (1) 1.71 A                                      (2) 2.0 A  
 (3) 2.31 A                                      (4) 1.33 A

---

Space For Rough Work

58. The input characteristics of a transistor in CE mode is the graph obtained by plotting

- (1)  $I_B$  against  $V_{BE}$  at constant  $V_{CE}$
- (2)  $I_B$  against  $V_{CE}$  at constant  $V_{BE}$
- (3)  $I_B$  against  $I_C$  at constant  $V_{CE}$
- (4)  $I_B$  against  $I_C$  at constant  $V_{BE}$

59. The given truth table is for

Input		Output
A	B	Y
0	0	1
0	1	1
1	0	1
1	1	0

- (1) AND gate
- (2) OR gate
- (3) NAND gate
- (4) NOR gate

60. The waves used for line-of-sight (LOS) communication is

- (1) ground waves
- (2) space waves
- (3) sound waves
- (4) sky waves

---

Space For Rough Work

A-I