

**Marking Scheme**  
**Mathematics (Term-I)**  
**Class-XII (Code-041)**

Q.N.	Correct Option	Hints / Solutions
1	d	$\sin\left(\frac{\pi}{3} - \left(-\frac{\pi}{6}\right)\right) = \sin\left(\frac{\pi}{2}\right) = 1$
2	b	$\lim_{x \rightarrow 0} \left(\frac{1 - \cos kx}{x \sin x}\right) = \frac{1}{2}$ $\Rightarrow \lim_{x \rightarrow 0} \left(\frac{2 \sin^2 \frac{kx}{2}}{x \sin x}\right) = \frac{1}{2}$ $\Rightarrow \lim_{x \rightarrow 0} 2 \left(\frac{k}{2}\right)^2 \left(\frac{\sin \frac{kx}{2}}{\frac{kx}{2}}\right)^2 \left(\frac{x}{\sin x}\right) = \frac{1}{2}$ $\Rightarrow k^2 = 1 \Rightarrow k = \pm 1 \text{ but } k < 0 \Rightarrow k = -1$
3	d	$A^2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
4	c	<p>As A is singular matrix</p> $\Rightarrow  A  = 0$ $\Rightarrow 2k^2 - 32 = 0 \Rightarrow k = \pm 4$
5	b	$f(x) = x^2 - 4x + 6$ $f'(x) = 2x - 4$ <p>let <math>f'(x) = 0 \Rightarrow x = 2</math></p> $\begin{array}{c} \longleftarrow \quad   \quad \longrightarrow \\ -\infty \quad (-) \quad 2 \quad (+) \quad \infty \end{array}$ <p>as <math>f'(x) &gt; 0 \quad \forall \quad x \in (2, \infty)</math>  <math>\Rightarrow f(x)</math> is Strictly increasing in <math>(2, \infty)</math></p>
6	d	<p>as <math> adj A  =  A ^{n-1}</math>, where n is order of matrix A</p> $= (-4)^2 = 16$
7	b	(1, 2)
8	a	$\left. \begin{array}{l} 2a + b = 4 \\ a - 2b = -3 \\ 5c - d = 11 \\ 4c + 3d = 24 \end{array} \right\} \Rightarrow \begin{array}{l} a = 1 \\ b = 2 \\ c = 3 \\ d = 4 \end{array}$ $\therefore a + b - c + 2d = 8$
9	a	$f(x) = x + \frac{1}{x}, x > 0 \Rightarrow f'(x) = 1 - \frac{1}{x^2} = \frac{x^2 - 1}{x^2}, x > 0$ <p>As normal to <math>f(x)</math> is <math>\perp</math> to given line</p> $\Rightarrow \left(\frac{x^2}{1-x^2}\right) \times \frac{3}{4} = -1 \quad (m_1 \cdot m_2 = -1)$ $\Rightarrow x^2 = 4 \Rightarrow x = \pm 2$ <p>But <math>x &gt; 0, \therefore x = 2</math></p> <p>Therefore point = <math>\left(2, \frac{5}{2}\right)</math></p>
10	d	$\sin(\tan^{-1} x) = \sin\left(\sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right)\right) = \frac{x}{\sqrt{1+x^2}}$
11	a	{1, 5, 9}
12	c	$e^x + e^y = e^{x+y}$ $\Rightarrow e^{-y} + e^{-x} = 1$ <p>Differentiating w.r.t. x:</p>

		$\Rightarrow -e^{-y} \frac{dy}{dx} - e^{-x} = 0 \Rightarrow \frac{dy}{dx} = -e^{y-x}$				
13	b	$3 \times 5$				
14	a	$y = 5 \cos x - 3 \sin x \Rightarrow \frac{dy}{dx} = -5 \sin x - 3 \cos x$ $\Rightarrow \frac{d^2y}{dx^2} = -5 \cos x + 3 \sin x = -y$				
15	c	$\text{adj } A = \begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix} \Rightarrow (\text{adj } A)' = \begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$				
16	c	$\frac{x^2}{9} + \frac{y^2}{16} = 1 \Rightarrow \frac{2x}{9} + \frac{2y}{16} \frac{dy}{dx} = 0$ $\Rightarrow \text{slope of normal} = \frac{-dx}{dy} = \frac{9y}{16x}$ As curve's tangent is parallel to y-axes $\Rightarrow \frac{9y}{16x} = 0 \Rightarrow y = 0$ and $x = \pm 3$ $\therefore \text{points} = (\pm 3, 0)$				
17	b	$ A  = -7$ $\therefore \sum_{i=1}^3 a_{i2} A_{i2} = a_{12} A_{12} + a_{22} A_{22} + a_{32} A_{32} =  A  = -7$				
18	d	$y = \log(\cos e^x)$ Differentiating wrt x: $\frac{dy}{dx} = \frac{1}{\cos(e^x)} \cdot (-\sin e^x) \cdot e^x$ (chain rule) $\Rightarrow \frac{dy}{dx} = -e^x \tan e^x$				
19	d	Z is maximum 180 at points C (15, 15) and D(0, 20). $\Rightarrow$ Z is maximum at every point on the line segment CD				
20	c	$f(x) = 2 \cos x + x, x \in [0, \frac{\pi}{2}]$ $f'(x) = -2 \sin x + 1$ Let $f'(x) = 0 \Rightarrow x = \frac{\pi}{6} \in [0, \frac{\pi}{2}]$ $f(0) = 2$ $f(\frac{\pi}{6}) = \frac{\pi}{6} + \sqrt{3}$ $f(\frac{\pi}{2}) = \frac{\pi}{2} \Rightarrow$ least value of $f(x)$ is $\frac{\pi}{2}$ at $x = \frac{\pi}{2}$				
<b>Section-B</b>						
21	d	<table border="0" style="width: 100%;"> <tr> <td style="width: 50%; vertical-align: top;"> <math>\text{let } f(x_1) = f(x_2) \forall x_1, x_2 \in R</math>  <math>\Rightarrow x_1^3 = x_2^3</math>  <math>\Rightarrow x_1 = x_2</math>  <math>\Rightarrow f</math> is one - one                 </td> <td style="width: 50%; vertical-align: top;"> <math>\text{let } f(x) = x^3 = y \forall y \in R</math>  <math>\Rightarrow x = y^{\frac{1}{3}}</math>                      every image <math>y \in R</math> has a unique pre image in <math>R</math>  <math>\Rightarrow f</math> is onto                 </td> </tr> <tr> <td colspan="2" style="text-align: center;"><math>\therefore f</math> is one-one and onto</td> </tr> </table>	$\text{let } f(x_1) = f(x_2) \forall x_1, x_2 \in R$ $\Rightarrow x_1^3 = x_2^3$ $\Rightarrow x_1 = x_2$ $\Rightarrow f$ is one - one	$\text{let } f(x) = x^3 = y \forall y \in R$ $\Rightarrow x = y^{\frac{1}{3}}$ every image $y \in R$ has a unique pre image in $R$ $\Rightarrow f$ is onto	$\therefore f$ is one-one and onto	
$\text{let } f(x_1) = f(x_2) \forall x_1, x_2 \in R$ $\Rightarrow x_1^3 = x_2^3$ $\Rightarrow x_1 = x_2$ $\Rightarrow f$ is one - one	$\text{let } f(x) = x^3 = y \forall y \in R$ $\Rightarrow x = y^{\frac{1}{3}}$ every image $y \in R$ has a unique pre image in $R$ $\Rightarrow f$ is onto					
$\therefore f$ is one-one and onto						
22	a	$x = a \sec \theta \Rightarrow \frac{dx}{d\theta} = a \tan \theta \sec \theta$ $y = b \tan \theta \Rightarrow \frac{dy}{d\theta} = b \sec^2 \theta$ $\therefore \frac{dy}{dx} = \frac{b}{a} \text{cosec } \theta$ $\Rightarrow \frac{d^2y}{dx^2} = \frac{-b}{a} \text{cosec } \theta \cdot \cot \theta \cdot \frac{d\theta}{dx} = \frac{-b}{a^2} \cot^3 \theta$ $\therefore \left. \frac{d^2y}{dx^2} \right _{\theta=\frac{\pi}{6}} = \frac{-3\sqrt{3}b}{a^2}$				
23	c	Z is minimum -24 at (0, 8)				
24	a	let $u = \sin^{-1}(2x\sqrt{1-x^2})$				



37	b	As every per-image $x \in A$ has a unique image $y \in B$ $\Rightarrow f$ is injective function
38	b	$ A  = 7, \text{adj}A = \begin{bmatrix} 2 & -2 \\ 1 & 3 \end{bmatrix}$ $\therefore 14A^{-1} = \begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$
39	b	$y = x^3 - 11x + 5 \Rightarrow \frac{dy}{dx} = 3x^2 - 11$ Slope of line $y = x - 11$ is 1 $\Rightarrow 3x^2 - 11 = 1 \Rightarrow x = \pm 2$ $\therefore$ point is (2, -9) as (-2, 19) does not satisfy given line
40	c	$A^2 = 3I$ $\Rightarrow \begin{bmatrix} \alpha^2 + \beta r & 0 \\ 0 & \beta r + \alpha^2 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} \Rightarrow 3 - \alpha^2 - \beta r = 0$
<b>Section C</b>		
41	a	As Z is maximum at (30, 30) and (0, 40) $\Rightarrow 30a + 30b = 40b \Rightarrow b - 3a = 0$
42	b	$y = mx + 1 \dots (1)$ and $y^2 = 4x \dots (2)$ Substituting (1) in (2): $(mx + 1)^2 = 4x$ $\Rightarrow m^2x^2 + (2m - 4)x + 1 = 0 \dots (3)$ As line is tangent to the curve $\Rightarrow$ line touches the curve at only one point $\Rightarrow (2m - 4)^2 - 4m^2 = 0 \Rightarrow m = 1$
43	c	Let $f(x) = [x(x - 1) + 1]^{\frac{1}{3}}, 0 \leq x \leq 1$ $f'(x) = \frac{2x-1}{3(x^2-x+1)^{\frac{2}{3}}}$ let $f'(x) = 0 \Rightarrow x = \frac{1}{2} \in [0, 1]$ $f(0) = 1, f\left(\frac{1}{2}\right) = \left(\frac{3}{4}\right)^{\frac{1}{3}}$ and $f(1) = 1$ $\therefore$ Maximum value of $f(x)$ is 1
44	b	Feasible region is bounded in the first quadrant
45	d	$ A  = 2 + 2\sin^2\theta$ As $-1 \leq \sin\theta \leq 1, \forall 0 \leq \theta \leq 2\pi$ $\Rightarrow 2 \leq 2 + 2\sin^2\theta \leq 4 \Rightarrow  A  \in [2, 4]$
46	d	Fuel cost = $k(\text{speed})^2$ $\Rightarrow 48 = k \cdot 16^2 \Rightarrow k = \frac{3}{16}$
47	b	Total cost of running train (let C) = $\frac{3}{16}v^2t + 1200t$ Distance covered = 500km $\Rightarrow$ time = $\frac{500}{v}$ hrs Total cost of running train 500 km = $\frac{3}{16}v^2\left(\frac{500}{v}\right) + 1200\left(\frac{500}{v}\right)$ $\Rightarrow C = \frac{375}{4}v + \frac{600000}{v}$
48	c	$\frac{dC}{dv} = \frac{375}{4} - \frac{600000}{v^2}$ Let $\frac{dC}{dv} = 0 \Rightarrow v = 80$ km/h
49	c	Fuel cost for running 500 km $\frac{375}{4}v = \frac{375}{4} \times 80 = \text{Rs. } 7500/-$
50	d	Total cost for running 500 km = $\frac{375}{4}v + \frac{600000}{v}$ $= \frac{375 \times 80}{4} + \frac{600000}{80} = \text{Rs. } 15000/-$