

JEE April 2019

Roll No.	
Candidate Name	
Application No	
Test Date	10/04/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

- Q.1** One mole of an ideal gas passes through a process where pressure and volume obey the relation $P = P_0 \left[1 - \frac{1}{2} \left(\frac{V_0}{V} \right)^2 \right]$. Here P_0 and V_0 are constants. Calculate the change in the temperature of the gas if its volume changes from V_0 to $2V_0$.

Options

1. $\frac{1}{2} \frac{P_0 V_0}{R}$
2. $\frac{5}{4} \frac{P_0 V_0}{R}$
3. $\frac{3}{4} \frac{P_0 V_0}{R}$
4. $\frac{1}{4} \frac{P_0 V_0}{R}$

Question Type : MCQ

Question ID : 41652914326

Option 1 ID : 41652956082

Option 2 ID : 41652956085

Option 3 ID : 41652956084

Option 4 ID : 41652956083


Status : Answered

Chosen Option : 1

- Q.2** The correct figure that shows, schematically, the wave pattern produced by superposition of two waves of frequencies 9 Hz and 11 Hz, is :

Options

- 1.
- 2.



Question Type : MCQ

Question ID : 41652914343

Option 1 ID : 41652956152

Option 2 ID : 41652956153

Option 3 ID : 41652956151

Option 4 ID : 41652956150

Status : Answered

Chosen Option : 4

Q.3 Light is incident normally on a completely absorbing surface with an energy flux of 25 Wcm^{-2} . If the surface has an area of 25 cm^2 , the momentum transferred to the surface in 40 min time duration will be :

- Options
1. $6.3 \times 10^{-4} \text{ Ns}$
 2. $1.4 \times 10^{-6} \text{ Ns}$
 3. $5.0 \times 10^{-3} \text{ Ns}$
 4. $3.5 \times 10^{-6} \text{ Ns}$

Question Type : MCQ

Question ID : 41652914336

Option 1 ID : 41652956125

Option 2 ID : 41652956122

Option 3 ID : 41652956124

Option 4 ID : 41652956123

Status : Not Answered

Chosen Option : --

Q.4 A solid sphere of mass M and radius R is divided into two unequal parts. The first part has a mass of $\frac{7M}{8}$ and is converted into a uniform disc of radius $2R$. The second part is converted into a uniform solid sphere. Let I_1 be the moment of inertia of the disc about its axis and I_2 be the moment of inertia of the new sphere about its axis. The ratio I_1/I_2 is given by :

- Options
1. 185
 2. 140

3. 285

4. 65

Question Type : MCQ

Question ID : 41652914321

Option 1 ID : 41652956065

Option 2 ID : 41652956062

Option 3 ID : 41652956064

Option 4 ID : 41652956063

Status : Answered

Chosen Option : 2

Q.5 A bullet of mass 20 g has an initial speed of 1 ms^{-1} , just before it starts penetrating a mud wall of thickness 20 cm. If the wall offers a mean resistance of $2.5 \times 10^{-2} \text{ N}$, the speed of the bullet after emerging from the other side of the wall is close to :

- Options
1. 0.1 ms^{-1}
 2. 0.7 ms^{-1}
 3. 0.3 ms^{-1}
 4. 0.4 ms^{-1}

Question Type : MCQ

Question ID : 41652914320

Option 1 ID : 41652956061

Option 2 ID : 41652956058

Option 3 ID : 41652956059

Option 4 ID : 41652956060

Status : Answered

Chosen Option : 4

Q.6 The time dependence of the position of a particle of mass $m=2$ is given by $\vec{r}(t) = 2t\hat{i} - 3t^2\hat{j}$. Its angular momentum, with respect to the origin, at time $t=2$ is :

- Options
1. $48(\hat{i} + \hat{j})$
 2. $36\hat{k}$
 3. $-34(\hat{k} - \hat{i})$
 4. $-48\hat{k}$

Question Type : MCQ

Question ID : 41652914318

Option 1 ID : 41652956053

Option 2 ID : 41652956050

Option 3 ID : 41652956052

Option 4 ID : 41652956051

Status : Answered

Chosen Option : 4

Q.7 A coil of self inductance 10 mH and resistance 0.1Ω is connected through a switch to a battery of internal resistance 0.9Ω . After the switch is closed, the time taken for the current to attain 80% of the saturation value is : [take $\ln 5 = 1.6$]

- Options
1. 0.324 s
 2. 0.103 s
 3. 0.002 s
 4. 0.016 s

Question Type : MCQ

Question ID : 41652914335

Option 1 ID : 41652956121

Option 2 ID : 41652956120

Option 3 ID : 41652956119

Option 4 ID : 41652956118

Status : Not Answered

Chosen Option : --

Q.8 In an experiment, brass and steel wires of length 1 m each with areas of cross section 1 mm^2 are used. The wires are connected in series and one end of the combined wire is connected to a rigid support and other end is subjected to elongation. The stress required to produce a net elongation of 0.2 mm is,

[Given, the Young's Modulus for steel and brass are, respectively, $120 \times 10^9 \text{ N/m}^2$ and $60 \times 10^9 \text{ N/m}^2$]

- Options
1. $1.2 \times 10^6 \text{ N/m}^2$
 2. $4.0 \times 10^6 \text{ N/m}^2$
 3. $1.8 \times 10^6 \text{ N/m}^2$
 4. $0.2 \times 10^6 \text{ N/m}^2$

Question Type : MCQ

Question ID : 41652914344

Option 1 ID : 41652956156

Option 2 ID : 41652956155

Option 3 ID : 41652956157

Option 4 ID : 41652956154

Status : Answered

Chosen Option : 3

Q.9

A source of sound S is moving with a velocity of 50 m/s towards a stationary observer. The observer measures the frequency of the source as 1000 Hz. What will be the apparent frequency of the source when it is moving away from the observer after crossing him ? (Take velocity of sound in air is 350 m/s)

- Options
1. 750 Hz
 2. 857 Hz
 3. 1143 Hz
 4. 807 Hz

Question Type : **MCQ**
Question ID : **41652914329**
Option 1 ID : **41652956096**
Option 2 ID : **41652956094**
Option 3 ID : **41652956095**
Option 4 ID : **41652956097**
Status : **Answered**
Chosen Option : **2**

Q.10 A 2 mW laser operates at a wavelength of 500 nm. The number of photons that will be emitted per second is :
[Given Planck's constant $h = 6.6 \times 10^{-34}$ Js, speed of light $c = 3.0 \times 10^8$ m/s]

- Options
1. 5×10^{15}
 2. 1.5×10^{16}
 3. 2×10^{16}
 4. 1×10^{16}

Question Type : **MCQ**
Question ID : **41652914339**
Option 1 ID : **41652956134**
Option 2 ID : **41652956136**
Option 3 ID : **41652956137**
Option 4 ID : **41652956135**
Status : **Answered**
Chosen Option : **1**

Q.11 A cubical block of side 0.5 m floats on water with 30% of its volume under water. What is the maximum weight that can be put on the block without fully submerging it under water ?
[Take, density of water = 10^3 kg/m³]

- Options
1. 46.3 kg
 2. 87.5 kg

3. 65.4 kg
4. 30.1 kg

Question Type : **MCQ**Question ID : **41652914345**Option 1 ID : **41652956160**Option 2 ID : **41652956159**Option 3 ID : **41652956161**Option 4 ID : **41652956158**Status : **Not Answered**

Chosen Option : --

Q.12 Two radioactive substances A and B have decay constants 5λ and λ respectively. At $t=0$, a sample has the same number of the two nuclei. The time taken for the ratio of the number of nuclei to become $\left(\frac{1}{e}\right)^2$ will be :

- Options
1. $1/2\lambda$
 2. $1/4\lambda$
 3. $1/\lambda$
 4. $2/\lambda$

Question Type : **MCQ**Question ID : **41652914341**Option 1 ID : **41652956144**Option 2 ID : **41652956145**Option 3 ID : **41652956142**Option 4 ID : **41652956143**Status : **Answered**Chosen Option : **4**

Q.13 Space between two concentric conducting spheres of radii a and b ($b > a$) is filled with a medium of resistivity ρ . The resistance between the two spheres will be :

- Options
1. $\frac{\rho}{4\pi} \left(\frac{1}{a} - \frac{1}{b} \right)$
 2. $\frac{\rho}{2\pi} \left(\frac{1}{a} - \frac{1}{b} \right)$
 3. $\frac{\rho}{2\pi} \left(\frac{1}{a} + \frac{1}{b} \right)$
 4. $\frac{\rho}{4\pi} \left(\frac{1}{a} + \frac{1}{b} \right)$

Question Type : **MCQ**Question ID : **41652914332**Option 1 ID : **41652956107**Option 2 ID : **41652956109**


Option 3 ID : **41652956108**
 Option 4 ID : **41652956106**
 Status : **Answered**
 Chosen Option : **4**

Q.14 In the formula $X = 5YZ^2$, X and Z have dimensions of capacitance and magnetic field, respectively. What are the dimensions of Y in SI units ?

- Options
1. $[M^{-3} L^{-2} T^8 A^4]$
 2. $[M^{-1} L^{-2} T^4 A^2]$
 3. $[M^{-2} L^0 T^{-4} A^{-2}]$
 4. $[M^{-2} L^{-2} T^6 A^3]$

Question Type : **MCQ**
 Question ID : **41652914316**
 Option 1 ID : **41652956043**
 Option 2 ID : **41652956042**
 Option 3 ID : **41652956045**
 Option 4 ID : **41652956044**
 Status : **Answered**
 Chosen Option : **3**

Q.15 The graph shows how the magnification m produced by a thin lens varies with image distance v. What is the focal length of the lens used ?



- Options
1. $\frac{b^2}{ac}$
 2. $\frac{b^2c}{a}$
 3. $\frac{a}{c}$
 4. $\frac{b}{c}$

Question Type : **MCQ**
 Question ID : **41652914337**
 Option 1 ID : **41652956126**
 Option 2 ID : **41652956128**
 Option 3 ID : **41652956127**
 Option 4 ID : **41652956129**
 Status : **Answered**
 Chosen Option : **4**

Q.16 A submarine experiences a pressure of 5.05×10^6 Pa at a depth of d_1 in a sea. When it goes further to a depth of d_2 , it experiences a pressure of 8.08×10^6 Pa. Then $d_2 - d_1$ is approximately (density of water = 10^3 kg/m³ and acceleration due to gravity = 10 ms^{-2}):

- Options
1. 300 m
 2. 400 m
 3. 600 m
 4. 500 m

Question Type : **MCQ**
Question ID : **41652914328**
Option 1 ID : **41652956090**
Option 2 ID : **41652956093**
Option 3 ID : **41652956091**
Option 4 ID : **41652956092**
Status : **Answered**
Chosen Option : **3**


Q.17 In a Young's double slit experiment, the ratio of the slit's width is 4 : 1. The ratio of the intensity of maxima to minima, close to the central fringe on the screen, will be :

- Options
1. 25 : 9
 2. 9 : 1
 3. 4 : 1
 4. $(\sqrt{3}+1)^4$: 16

Question Type : **MCQ**
Question ID : **41652914338**
Option 1 ID : **41652956133**
Option 2 ID : **41652956131**
Option 3 ID : **41652956130**
Option 4 ID : **41652956132**
Status : **Answered**
Chosen Option : **3**

Q.18

A metal coin of mass 5 g and radius 1 cm is fixed to a thin stick AB of negligible mass as shown in the figure. The system is initially at rest. The constant torque, that will make the system rotate about AB at 25 rotations per second in 5 s, is close to :



- Options
1. $4.0 \times 10^{-6} \text{ Nm}$
 2. $1.6 \times 10^{-5} \text{ Nm}$
 3. $7.9 \times 10^{-6} \text{ Nm}$
 4. $2.0 \times 10^{-5} \text{ Nm}$

Question Type : **MCQ**

Question ID : **41652914322**

Option 1 ID : **41652956066**

Option 2 ID : **41652956068**

Option 3 ID : **41652956067**

Option 4 ID : **41652956069**

Status : **Answered**

Chosen Option : **4**

Q.19 In free space, a particle A of charge $1 \mu\text{C}$ is held fixed at a point P. Another particle B of the same charge and mass $4 \mu\text{g}$ is kept at a distance of 1 mm from P. If B is released, then its velocity at a distance of 9 mm from P is :

$$\left[\text{Take } \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2} \right]$$

- Options
1. 1.0 m/s
 2. $3.0 \times 10^4 \text{ m/s}$
 3. $2.0 \times 10^3 \text{ m/s}$
 4. $1.5 \times 10^2 \text{ m/s}$

Question Type : **MCQ**

Question ID : **41652914330**

Option 1 ID : **41652956101**

Option 2 ID : **41652956098**

Option 3 ID : **41652956099**

Option 4 ID : **41652956100**

Status : **Not Answered**

Chosen Option : **--**

Q.20

The elastic limit of brass is 379 MPa. What should be the minimum diameter of a brass rod if it is to support a 400 N load without exceeding its elastic limit ?

- Options
1. 1.00 mm
 2. 1.16 mm
 3. 0.90 mm
 4. 1.36 mm

Question Type : **MCQ**

Question ID : **41652914324**

Option 1 ID : **41652956074**

Option 2 ID : **41652956076**

Option 3 ID : **41652956075**

Option 4 ID : **41652956077**

Status : **Answered**

Chosen Option : **3**

Q.21 A spaceship orbits around a planet at a height of 20 km from its surface. Assuming that only gravitational field of the planet acts on the spaceship, what will be the number of complete revolutions made by the spaceship in 24 hours around the planet ?

[Given : Mass of planet = 8×10^{22} kg,

Radius of planet = 2×10^6 m,

Gravitational constant $G = 6.67 \times 10^{-11}$ Nm^2/kg^2]

- Options
1. 9
 2. 17
 3. 13
 4. 11

Question Type : **MCQ**

Question ID : **41652914323**

Option 1 ID : **41652956072**

Option 2 ID : **41652956071**

Option 3 ID : **41652956070**

Option 4 ID : **41652956073**

Status : **Answered**

Chosen Option : **4**

Q.22

In Li^{++} , electron in first Bohr orbit is excited to a level by a radiation of wavelength λ . When the ion gets deexcited to the ground state in all possible ways (including intermediate emissions), a total of six spectral lines are observed. What is the value of λ ?

(Given : $h = 6.63 \times 10^{-34} \text{ Js}$;

$c = 3 \times 10^8 \text{ ms}^{-1}$)

- Options
1. 11.4 nm
 2. 9.4 nm
 3. 12.3 nm
 4. 10.8 nm

Question Type : MCQ

Question ID : 41652914340

Option 1 ID : 41652956140

Option 2 ID : 41652956138

Option 3 ID : 41652956141


Option 4 ID : 41652956139

Status : Not Answered

Chosen Option : --

Q.23

A simple pendulum of length L is placed between the plates of a parallel plate capacitor having electric field E , as shown in figure. Its bob has mass m and charge q . The time period of the pendulum is given by :



- Options
1. $2\pi \sqrt{\frac{L}{\left(g + \frac{qE}{m}\right)}}$
 2. $2\pi \sqrt{\frac{L}{\sqrt{g^2 - \frac{q^2 E^2}{m^2}}}}$
 3. $2\pi \sqrt{\frac{L}{\left(g - \frac{qE}{m}\right)}}$

4.
$$2\pi \sqrt{\frac{L}{\sqrt{g^2 + \left(\frac{qE}{m}\right)^2}}}$$

Question Type : MCQ

Question ID : 41652914331

Option 1 ID : 41652956102

Option 2 ID : 41652956105

Option 3 ID : 41652956103

Option 4 ID : 41652956104

Status : Answered

Chosen Option : 3

Q.24 The magnitude of the magnetic field at the center of an equilateral triangular loop of side 1 m which is carrying a current of 10 A is :

[Take $\mu_0 = 4\pi \times 10^{-7} \text{ NA}^{-2}$]

- Options
1. $18 \mu\text{T}$
 2. $9 \mu\text{T}$
 3. $3 \mu\text{T}$
 4. $1 \mu\text{T}$

Question Type : MCQ

Question ID : 41652914333

Option 1 ID : 41652956113

Option 2 ID : 41652956112


Option 3 ID : 41652956111

Option 4 ID : 41652956110

Status : Answered

Chosen Option : 2

Q.25 The figure represents a voltage regulator circuit using a Zener diode. The breakdown voltage of the Zener diode is 6 V and the load resistance is, $R_L = 4 \text{ k}\Omega$. The series resistance of the circuit is $R_i = 1 \text{ k}\Omega$. If the battery voltage V_B varies from 8 V to 16 V, what are the minimum and maximum values of the current through Zener diode ?



- Options
1. 0.5 mA; 6 mA
 2. 1 mA; 8.5 mA
 3. 0.5 mA; 8.5 mA

4. 1.5 mA; 8.5 mA

Question Type : MCQ

Question ID : 41652914342

Option 1 ID : 41652956146

Option 2 ID : 41652956149

Option 3 ID : 41652956148

Option 4 ID : 41652956147

Status : Not Answered

Chosen Option : --

Q.26 When heat Q is supplied to a diatomic gas of rigid molecules, at constant volume its temperature increases by ΔT . The heat required to produce the same change in temperature, at a constant pressure is :

Options

1. $\frac{2}{3}Q$
2. $\frac{5}{3}Q$
3. $\frac{7}{5}Q$
4. $\frac{3}{2}Q$

Question Type : MCQ

Question ID : 41652914327

Option 1 ID : 41652956088

Option 2 ID : 41652956086

Option 3 ID : 41652956087


Option 4 ID : 41652956089

Status : Not Answered

Chosen Option : --

Q.27 Two blocks A and B of masses $m_A = 1$ kg and $m_B = 3$ kg are kept on the table as shown in figure. The coefficient of friction between A and B is 0.2 and between B and the surface of the table is also 0.2. The maximum force F that can be applied on B horizontally, so that the block A does not slide over the block B is :

[Take $g = 10 \text{ m/s}^2$]



Options

1. 8 N
2. 16 N
3. 40 N
4. 12 N

Question Type : **MCQ**Question ID : **41652914319**Option 1 ID : **41652956054**Option 2 ID : **41652956056**Option 3 ID : **41652956057**Option 4 ID : **41652956055**Status : **Answered**Chosen Option : **2**

Q.28 Water from a tap emerges vertically downwards with an initial speed of 1.0 ms^{-1} . The cross-sectional area of the tap is 10^{-4} m^2 . Assume that the pressure is constant throughout the stream of water and that the flow is streamlined. The cross-sectional area of the stream, 0.15 m below the tap would be :
(Take $g = 10 \text{ ms}^{-2}$)

- Options
1. $2 \times 10^{-5} \text{ m}^2$
 2. $5 \times 10^{-5} \text{ m}^2$
 3. $5 \times 10^{-4} \text{ m}^2$
 4. $1 \times 10^{-5} \text{ m}^2$

Question Type : **MCQ**Question ID : **41652914325**Option 1 ID : **41652956079**Option 2 ID : **41652956081**Option 3 ID : **41652956080**Option 4 ID : **41652956078**Status : **Not Answered**Chosen Option : **--**

Q.29 A square loop is carrying a steady current I and the magnitude of its magnetic dipole moment is m . If this square loop is changed to a circular loop and it carries the same current, the magnitude of the magnetic dipole moment of circular loop will be :

- Options
1. $\frac{m}{\pi}$
 2. $\frac{3m}{\pi}$
 3. $\frac{2m}{\pi}$
 4. $\frac{4m}{\pi}$

Question Type : **MCQ**Question ID : **41652914334**Option 1 ID : **41652956114**Option 2 ID : **41652956116**

Option 3 ID : 41652956115

Option 4 ID : 41652956117


Status : Answered

Chosen Option : 3

Q.30

A plane is inclined at an angle $\alpha = 30^\circ$ with respect to the horizontal. A particle is projected with a speed $u = 2 \text{ ms}^{-1}$ from the base of the plane, making an angle $\theta = 15^\circ$ with respect to the plane as shown in the figure. The distance from the base, at which the particle hits the plane is close to :

(Take $g = 10 \text{ ms}^{-2}$)



- Options
1. 20 cm
 2. 18 cm
 3. 26 cm
 4. 14 cm

Question Type : MCQ

Question ID : 41652914317

Option 1 ID : 41652956048

Option 2 ID : 41652956047

Option 3 ID : 41652956049

Option 4 ID : 41652956046

Status : Not Answered

Chosen Option : --

Section : Chemistry

Q.1 Number of stereo centers present in linear and cyclic structures of glucose are respectively :

- Options
1. 5 & 4
 2. 4 & 4
 3. 5 & 5
 4. 4 & 5

Question Type : MCQ

Question ID : 41652914349

Option 1 ID : 41652956176

Option 2 ID : 41652956174

Option 3 ID : 41652956177

Option 4 ID : 41652956175

Status : Answered

Chosen Option : 4


Q.2

The pH of a 0.02 M NH_4Cl solution will be
[given $K_b(\text{NH}_4\text{OH}) = 10^{-5}$ and $\log 2 = 0.301$]

- Options
1. 2.65
 2. 4.35
 3. 4.65
 4. 5.35

Question Type : **MCQ**
 Question ID : **41652914372**
 Option 1 ID : **41652956267**
 Option 2 ID : **41652956269**
 Option 3 ID : **41652956266**
 Option 4 ID : **41652956268**
 Status : **Answered**
 Chosen Option : **2**

Q.3 The major product 'Y' in the following reaction is :



Options

- 1.
- 2.
- 3.
- 4.

Question Type : **MCQ**
 Question ID : **41652914352**
 Option 1 ID : **41652956189**
 Option 2 ID : **41652956187**
 Option 3 ID : **41652956188**
 Option 4 ID : **41652956186**
 Status : **Answered**
 Chosen Option : **3**

Q.4 Which of the following is NOT a correct method of the preparation of benzylamine from cyanobenzene ?

Options

1. H_2/Ni
2. (i) LiAlH_4 (ii) H_3O^+
3. (i) $\text{SnCl}_2 + \text{HCl}(\text{gas})$ (ii) NaBH_4
4. (i) $\text{HCl}/\text{H}_2\text{O}$ (ii) NaBH_4

Question Type : MCQ

Question ID : 41652914355

Option 1 ID : 41652956200

Option 2 ID : 41652956199

Option 3 ID : 41652956201

Option 4 ID : 41652956198

Status : Answered

Chosen Option : 4

Q.5 The highest possible oxidation states of uranium and plutonium, respectively, are :

Options

1. 6 and 7
2. 6 and 4
3. 7 and 6
4. 4 and 6

Question Type : MCQ

Question ID : 41652914362

Option 1 ID : 41652956227

Option 2 ID : 41652956228

Option 3 ID : 41652956226

Option 4 ID : 41652956229

Status : Answered

Chosen Option : 2

Q.6 A hydrated solid X on heating initially gives a monohydrated compound Y. Y upon heating above 373 K leads to an anhydrous white powder Z. X and Z, respectively, are :

Options

1. Washing soda and soda ash.
2. Baking soda and dead burnt plaster.
3. Washing soda and dead burnt plaster.
4. Baking soda and soda ash.

Question Type : MCQ

Question ID : 41652914359

Option 1 ID : 41652956216

Option 2 ID : 41652956217

Option 3 ID : 41652956215

Option 4 ID : 41652956214

Status : Marked For Review

Chosen Option : 3

Q.7

The correct order of the first ionization enthalpies is :

- Options
1. $\text{Ti} < \text{Mn} < \text{Zn} < \text{Ni}$
 2. $\text{Ti} < \text{Mn} < \text{Ni} < \text{Zn}$
 3. $\text{Mn} < \text{Ti} < \text{Zn} < \text{Ni}$
 4. $\text{Zn} < \text{Ni} < \text{Mn} < \text{Ti}$

Question Type : **MCQ**

Question ID : **41652914356**

Option 1 ID : **41652956204**

Option 2 ID : **41652956202**

Option 3 ID : **41652956203**

Option 4 ID : **41652956205**

Status : **Answered**

Chosen Option : **1**

Q.8 The correct statements among (a) to (d) are :

- (a) saline hydrides produce H_2 gas when reacted with H_2O .
- (b) reaction of LiAlH_4 with BF_3 leads to B_2H_6 .
- (c) PH_3 and CH_4 are electron - rich and electron - precise hydrides, respectively.
- (d) HF and CH_4 are called as molecular hydrides.

- Options
1. (a), (b), (c) and (d).
 2. (c) and (d) only.
 3. (a), (c) and (d) only.
 4. (a), (b) and (c) only.

Question Type : **MCQ**

Question ID : **41652914358**

Option 1 ID : **41652956213**

Option 2 ID : **41652956210**

Option 3 ID : **41652956211**

Option 4 ID : **41652956212**

Status : **Answered**

Chosen Option : **3**

Q.9 The ratio of the shortest wavelength of two spectral series of hydrogen spectrum is found to be about 9. The spectral series are :

- Options
1. Lyman and Paschen
 2. Balmer and Brackett

3. Brackett and Pfund
4. Paschen and Pfund

Question Type : **MCQ**Question ID : **41652914368**Option 1 ID : **41652956251**Option 2 ID : **41652956250**Option 3 ID : **41652956253**Option 4 ID : **41652956252**Status : **Not Attempted and Marked For Review**

Chosen Option : --

Q.10 The correct statement is :

- Options
1. aniline is a froth stabilizer.
 2. zincite is a carbonate ore.
 3. sodium cyanide cannot be used in the metallurgy of silver.
 4. zone refining process is used for the refining of titanium.

Question Type : **MCQ**Question ID : **41652914357**Option 1 ID : **41652956206**Option 2 ID : **41652956208**Option 3 ID : **41652956207**Option 4 ID : **41652956209**Status : **Answered**Chosen Option : **3**

Q.11 The increasing order of nucleophilicity of the following nucleophiles is :

- (a) CH_3CO_2^-
- (b) H_2O
- (c) CH_3SO_3^-
- (d) OH^-

- Options
1. (a) < (d) < (c) < (b)
 2. (b) < (c) < (d) < (a)
 3. (d) < (a) < (c) < (b)
 4. (b) < (c) < (a) < (d)

Question Type : **MCQ**Question ID : **41652914354**Option 1 ID : **41652956195**Option 2 ID : **41652956194**Option 3 ID : **41652956197**Option 4 ID : **41652956196**Status : **Answered**Chosen Option : **4**

Q.12

The correct option among the following is :

- Options
1. Colloidal medicines are more effective because they have small surface area.
 2. Addition of alum to water makes it unfit for drinking.
 3. Colloidal particles in lyophobic sols can be precipitated by electrophoresis.
 4. Brownian motion in colloidal solution is faster if the viscosity of the solution is very high.

Question Type : **MCQ**
Question ID : **41652914375**
Option 1 ID : **41652956279**
Option 2 ID : **41652956278**
Option 3 ID : **41652956280**
Option 4 ID : **41652956281**
Status : **Answered**
Chosen Option : **3**

Q.13 Which of these factors does not govern the stability of a conformation in acyclic compounds ?

- Options
1. Steric interactions
 2. Torsional strain
 3. Electrostatic forces of interaction
 4. Angle strain

Question Type : **MCQ**
Question ID : **41652914347**
Option 1 ID : **41652956166**
Option 2 ID : **41652956168**
Option 3 ID : **41652956169**
Option 4 ID : **41652956167**
Status : **Answered**
Chosen Option : **1**

Q.14 In chromatography, which of the following statements is INCORRECT for R_f ?

- Options
1. R_f value depends on the type of chromatography.
 2. The value of R_f can not be more than one.
 3. Higher R_f value means higher adsorption.

4. R_f value is dependent on the mobile phase.

Question Type : MCQ

Question ID : 41652914348

Option 1 ID : 41652956171

Option 2 ID : 41652956172

Option 3 ID : 41652956170

Option 4 ID : 41652956173

Status : Not Answered

Chosen Option : --

- Q.15 The crystal field stabilization energy (CFSE) of $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_2$ and $\text{K}_2[\text{NiCl}_4]$, respectively, are :

- Options
1. $-0.6\Delta_o$ and $-0.8\Delta_t$
 2. $-0.4\Delta_o$ and $-0.8\Delta_t$
 3. $-2.4\Delta_o$ and $-1.2\Delta_t$
 4. $-0.4\Delta_o$ and $-1.2\Delta_t$

Question Type : MCQ

Question ID : 41652914363

Option 1 ID : 41652956230

Option 2 ID : 41652956231

Option 3 ID : 41652956232

Option 4 ID : 41652956233

Status : Answered

Chosen Option : 2

- Q.16 The difference between ΔH and ΔU ($\Delta H - \Delta U$), when the combustion of one mole of heptane(l) is carried out at a temperature T, is equal to :

- Options
1. $-4 RT$
 2. $-3 RT$
 3. $4 RT$
 4. $3 RT$

Question Type : MCQ

Question ID : 41652914369

Option 1 ID : 41652956256

Option 2 ID : 41652956257

Option 3 ID : 41652956254





Option 4 ID : 41652956255

Status : Answered

Chosen Option : 4

- Q.17 Which one of the following graphs between molar conductivity (Λ_m) versus $\sqrt{C}$ is correct ?

Options



Question Type : MCQ

Question ID : 41652914373

Option 1 ID : 41652956270

Option 2 ID : 41652956273

Option 3 ID : 41652956271

Option 4 ID : 41652956272

Status : Not Answered

Chosen Option : --

Q.18 The correct match between Item - I and Item - II is :


	Item - I		Item - II
(a)	High density polythene	(I)	Peroxide catalyst
(b)	Polyacrylonitrile	(II)	Condensation at high temperature & pressure
(c)	Novolac	(III)	Ziegler-Natta Catalyst
(d)	Nylon 6	(IV)	Acid or base catalyst

Options
1. (a) → (IV), (b) → (II), (c) → (I),
(d) → (III)

2. (a) $\rightarrow$ (II), (b) $\rightarrow$ (IV), (c) $\rightarrow$ (I),
(d) $\rightarrow$ (III)
3. (a) $\rightarrow$ (III), (b) $\rightarrow$ (I), (c) $\rightarrow$ (II),
(d) $\rightarrow$ (IV)
4. (a) $\rightarrow$ (III), (b) $\rightarrow$ (I), (c) $\rightarrow$ (IV),
(d) $\rightarrow$ (II)

Question Type : **MCQ**Question ID : **41652914350**Option 1 ID : **41652956178**Option 2 ID : **41652956179**Option 3 ID : **41652956180**Option 4 ID : **41652956181**Status : **Answered**Chosen Option : **3**


Q.19 Points I, II and III in the following plot respectively correspond to
(V_{mp} : most probable velocity)



- Options
1. V_{mp} of N_2 (300 K); V_{mp} of O_2 (400 K);
 V_{mp} of H_2 (300 K)
 2. V_{mp} of O_2 (400 K); V_{mp} of N_2 (300 K);
 V_{mp} of H_2 (300 K)
 3. V_{mp} of N_2 (300 K); V_{mp} of H_2 (300 K);
 V_{mp} of O_2 (400 K)
 4. V_{mp} of H_2 (300 K); V_{mp} of N_2 (300 K);
 V_{mp} of O_2 (400 K)

Question Type : **MCQ**Question ID : **41652914367**Option 1 ID : **41652956248**Option 2 ID : **41652956247**Option 3 ID : **41652956249**Option 4 ID : **41652956246**Status : **Answered**Chosen Option : **2****Q.20**

The major product obtained in the given reaction is :



Options

- 1.
- 2.
- 3.
- 4.

Question Type : MCQ

Question ID : 41652914353

Option 1 ID : 41652956191

Option 2 ID : 41652956193

Option 3 ID : 41652956192

Option 4 ID : 41652956190

Status : Answered

Chosen Option : 3

Q.21 The minimum amount of $\text{O}_2(\text{g})$ consumed per gram of reactant is for the reaction :

(Given atomic mass : Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)

Options

1. $4 \text{ Fe}(\text{s}) + 3 \text{ O}_2(\text{g}) \rightarrow 2 \text{ Fe}_2\text{O}_3(\text{s})$
2. $\text{P}_4(\text{s}) + 5 \text{ O}_2(\text{g}) \rightarrow \text{P}_4\text{O}_{10}(\text{s})$
3. $\text{C}_3\text{H}_8(\text{g}) + 5 \text{ O}_2(\text{g}) \rightarrow 3 \text{ CO}_2(\text{g}) + 4 \text{ H}_2\text{O}(\text{l})$
4. $2 \text{ Mg}(\text{s}) + \text{O}_2(\text{g}) \rightarrow 2 \text{ MgO}(\text{s})$

Question Type : MCQ

Question ID : 41652914366

Option 1 ID : 41652956242

Option 2 ID : 41652956244

Option 3 ID : 41652956245


Option 4 ID : 41652956243

Status : Answered

Chosen Option : 1

Q.22

For the reaction,



$$\Delta H = -57.2 \text{ kJ mol}^{-1} \text{ and}$$

$$K_c = 1.7 \times 10^{16}.$$

Which of the following statement is INCORRECT?

- Options
1. The equilibrium constant is large suggestive of reaction going to completion and so no catalyst is required.
 2. The equilibrium will shift in forward direction as the pressure increases.
 3. The equilibrium constant decreases as the temperature increases.
 4. The addition of inert gas at constant volume will not affect the equilibrium constant.

Question Type : **MCQ**

Question ID : **41652914371**

Option 1 ID : **41652956265**

Option 2 ID : **41652956263**

Option 3 ID : **41652956262**

Option 4 ID : **41652956264**

Status : **Answered**

Chosen Option : **3**

- Q.23** 1 g of a non-volatile non-electrolyte solute is dissolved in 100 g of two different solvents A and B whose ebullioscopic constants are in the ratio of 1 : 5. The ratio of the elevation in their boiling points,

$$\frac{\Delta T_b(\text{A})}{\Delta T_b(\text{B})}, \text{ is :}$$

- Options
1. 5 : 1
 2. 10 : 1
 3. 1 : 5
 4. 1 : 0.2

Question Type : **MCQ**

Question ID : **41652914370**

Option 1 ID : **41652956259**

Option 2 ID : **41652956261**

Option 3 ID : **41652956260**

Option 4 ID : **41652956258**

Status : **Not Answered**

Chosen Option : **--**

Q.24

For the reaction of H_2 with I_2 , the rate constant is $2.5 \times 10^{-4} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ at 327°C and $1.0 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ at 527°C . The activation energy for the reaction, in kJ mol^{-1} is :

($R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$)

Options 1. 166

2. 150

3. 72

4. 59

Question Type : MCQ

Question ID : 41652914374

Option 1 ID : 41652956277

Option 2 ID : 41652956276

Option 3 ID : 41652956275

Option 4 ID : 41652956274

Status : Not Attempted and Marked For Review

Chosen Option : --

Q.25 The INCORRECT statement is :

Options the gemstone, ruby, has Cr^{3+} ions

1. occupying the octahedral sites of beryl.

2. the spin-only magnetic moment of $[\text{Ni}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$ is 2.83 BM.

3. the color of $[\text{CoCl}(\text{NH}_3)_5]^{2+}$ is violet as it absorbs the yellow light.

4. the spin-only magnetic moments of $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ and $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$ are nearly similar.

Question Type : MCQ

Question ID : 41652914364

Option 1 ID : 41652956237

Option 2 ID : 41652956234

Option 3 ID : 41652956236





Option 4 ID : 41652956235

Status : Answered

Chosen Option : 2

Q.26 Compound A ($\text{C}_9\text{H}_{10}\text{O}$) shows positive iodoform test. Oxidation of A with KMnO_4/KOH gives acid B ($\text{C}_8\text{H}_6\text{O}_4$). Anhydride of B is used for the preparation of phenolphthalein. Compound A is :

Options



Question Type : **MCQ**

Question ID : **41652914351**

Option 1 ID : **41652956182**

Option 2 ID : **41652956185**

Option 3 ID : **41652956184**

Option 4 ID : **41652956183**

Status : **Answered**

Chosen Option : **4**

Q.27 The number of pentagons in C_{60} and trigons (triangles) in white phosphorus, respectively, are :

- Options
1. 20 and 3
 2. 12 and 4
 3. 12 and 3
 4. 20 and 4

Question Type : **MCQ**

Question ID : **41652914361**

Option 1 ID : **41652956224**

Option 2 ID : **41652956222**

Option 3 ID : **41652956223**

Option 4 ID : **41652956225**

Status : **Not Answered**

Chosen Option : **--**

Q.28 Air pollution that occurs in sunlight is :

- Options
1. reducing smog
 2. acid rain

3. oxidising smog
4. fog


Question Type : MCQ
Question ID : 41652914365
Option 1 ID : 41652956240
Option 2 ID : 41652956238
Option 3 ID : 41652956241
Option 4 ID : 41652956239
Status : Answered
Chosen Option : 3

Q.29 The noble gas that does NOT occur in the atmosphere is :



- Options
1. He
 2. Kr
 3. Ne
 4. Ra


Question Type : MCQ
Question ID : 41652914360
Option 1 ID : 41652956218
Option 2 ID : 41652956220
Option 3 ID : 41652956219
Option 4 ID : 41652956221
Status : Answered
Chosen Option : 4

Q.30 The major product 'Y' in the following reaction is :



Options

1. 
2. 

Question Type : **MCQ**Question ID : **41652914346**Option 1 ID : **41652956164**Option 2 ID : **41652956163**Option 3 ID : **41652956165**Option 4 ID : **41652956162**Status : **Answered**Chosen Option : **2**Section : **Mathematics**

Q.1 If $5x + 9 = 0$ is the directrix of the hyperbola $16x^2 - 9y^2 = 144$, then its corresponding focus is :

Options 1. $(5, 0)$ 2. $\left(-\frac{5}{3}, 0\right)$ 3. $\left(\frac{5}{3}, 0\right)$ 4. $(-5, 0)$ Question Type : **MCQ**Question ID : **41652914397**Option 1 ID : **41652956367**Option 2 ID : **41652956368**Option 3 ID : **41652956369**Option 4 ID : **41652956366**Status : **Answered**Chosen Option : **4**

Q.2 The number of real roots of the equation

$$5 + |2^x - 1| = 2^x(2^x - 2) \text{ is :}$$

Options 1. **3**2. **2**3. **4**4. **1**Question Type : **MCQ**Question ID : **41652914378**

Option 1 ID : **41652956293**
 Option 2 ID : **41652956292**
 Option 3 ID : **41652956290**
 Option 4 ID : **41652956291**
 Status : **Answered**
 Chosen Option : **4**

Q.3 The tangent and normal to the ellipse $3x^2 + 5y^2 = 32$ at the point $P(2, 2)$ meet the x -axis at Q and R , respectively. Then the area (in sq. units) of the triangle PQR is :

- Options
1. $\frac{34}{15}$
 2. $\frac{14}{3}$
 3. $\frac{16}{3}$
 4. $\frac{68}{15}$

Question Type : **MCQ**
 Question ID : **41652914396**
 Option 1 ID : **41652956362**
 Option 2 ID : **41652956364**
 Option 3 ID : **41652956363**
 Option 4 ID : **41652956365**
 Status : **Answered**
 Chosen Option : **4**

Q.4 Let $f(x) = \log_e(\sin x)$, $(0 < x < \pi)$ and $g(x) = \sin^{-1}(e^{-x})$, $(x \geq 0)$. If α is a positive real number such that $a = (f \circ g)'(\alpha)$ and $b = (f \circ g)(\alpha)$, then :

- Options
1. $a\alpha^2 + b\alpha + a = 0$
 2. $a\alpha^2 - b\alpha - a = 1$
 3. $a\alpha^2 - b\alpha - a = 0$
 4. $a\alpha^2 + b\alpha - a = -2\alpha^2$

Question Type : **MCQ**
 Question ID : **41652914376**
 Option 1 ID : **41652956283**
 Option 2 ID : **41652956285**
 Option 3 ID : **41652956282**
 Option 4 ID : **41652956284**
 Status : **Answered**
 Chosen Option : **3**

Q.5

Suppose that 20 pillars of the same height have been erected along the boundary of a circular stadium. If the top of each pillar has been connected by beams with the top of all its non-adjacent pillars, then the total number of beams is :

- Options
1. 170
 2. 180
 3. 210
 4. 190

Question Type : **MCQ**

Question ID : **41652914381**

Option 1 ID : **41652956302**

Option 2 ID : **41652956304**

Option 3 ID : **41652956305**

Option 4 ID : **41652956303**

Status : **Answered**

Chosen Option : **1**

Q.6 If the tangent to the curve $y = \frac{x}{x^2 - 3}$, $x \in \mathbb{R}$, ($x \neq \pm\sqrt{3}$), at a point $(\alpha, \beta) \neq (0, 0)$ on it is parallel to the line $2x + 6y - 11 = 0$, then :

- Options
1. $|6\alpha + 2\beta| = 19$
 2. $|6\alpha + 2\beta| = 9$
 3. $|2\alpha + 6\beta| = 19$
 4. $|2\alpha + 6\beta| = 11$

Question Type : **MCQ**

Question ID : **41652914388**

Option 1 ID : **41652956330**

Option 2 ID : **41652956332**

Option 3 ID : **41652956333**

Option 4 ID : **41652956331**

Status : **Answered**

Chosen Option : **3**

Q.7 If the line $ax + y = c$, touches both the curves $x^2 + y^2 = 1$ and $y^2 = 4\sqrt{2}x$, then $|c|$ is equal to :

- Options
1. 2
 2. $\frac{1}{\sqrt{2}}$
 3. $\frac{1}{2}$

4. $\sqrt{2}$

Question Type : MCQ

Question ID : 41652914395

Option 1 ID : 41652956361

Option 2 ID : 41652956359

Option 3 ID : 41652956360

Option 4 ID : 41652956358

Status : Answered

Chosen Option : 4

Q.8

The sum $1 + \frac{1^3+2^3}{1+2} + \frac{1^3+2^3+3^3}{1+2+3} + \dots$

$+ \frac{1^3+2^3+3^3+\dots+15^3}{1+2+3+\dots+15} - \frac{1}{2}(1+2+3+\dots+15)$

is equal to :

Options

1. 620

2. 1240

3. 1860

4. 660

Question Type : MCQ

Question ID : 41652914384

Option 1 ID : 41652956314

Option 2 ID : 41652956316

Option 3 ID : 41652956317

Option 4 ID : 41652956315

Status : Not Attempted and Marked For Review

Chosen Option : --

Q.9

If z and w are two complex numbers such

that $|zw| = 1$ and $\arg(z) - \arg(w) = \frac{\pi}{2}$,

then :

Options

1. $\bar{z}w = i$ 2. $\bar{z}w = \frac{-1+i}{\sqrt{2}}$ 3. $\bar{z}w = -i$ 4. $\bar{z}w = \frac{1-i}{\sqrt{2}}$

Question Type : MCQ

Question ID : 41652914377

Option 1 ID : 41652956288

Option 2 ID : 41652956287

Option 3 ID : 41652956286

Option 4 ID : 41652956289

Status : Not Answered

Chosen Option : --

Q.10

Let λ be a real number for which the system of linear equations

$$x + y + z = 6$$

$$4x + \lambda y - \lambda z = \lambda - 2$$

$$3x + 2y - 4z = -5$$

has infinitely many solutions. Then λ is a root of the quadratic equation :

- Options
1. $\lambda^2 + 3\lambda - 4 = 0$
 2. $\lambda^2 - 3\lambda - 4 = 0$
 3. $\lambda^2 + \lambda - 6 = 0$
 4. $\lambda^2 - \lambda - 6 = 0$

Question Type : **MCQ**

Question ID : **41652914380**

Option 1 ID : **41652956298**

Option 2 ID : **41652956299**

Option 3 ID : **41652956300**

Option 4 ID : **41652956301**

Status : **Answered**

Chosen Option : **4**

Q.11 If $\cos^{-1} x - \cos^{-1} \frac{y}{2} = \alpha$, where $-1 \leq x \leq 1$,
 $-2 \leq y \leq 2$, $x \leq \frac{y}{2}$, then for all x, y ,
 $4x^2 - 4xy \cos \alpha + y^2$ is equal to :

- Options
1. $4 \sin^2 \alpha$
 2. $2 \sin^2 \alpha$
 3. $4 \sin^2 \alpha - 2x^2 y^2$
 4. $4 \cos^2 \alpha + 2x^2 y^2$

Question Type : **MCQ**

Question ID : **41652914403**

Option 1 ID : **41652956392**

Option 2 ID : **41652956390**

Option 3 ID : **41652956391**

Option 4 ID : **41652956393**

Status : **Answered**

Chosen Option : **3**

Q.12 The distance of the point having position vector $-\hat{i} + 2\hat{j} + 6\hat{k}$ from the straight line passing through the point $(2, 3, -4)$ and parallel to the vector, $6\hat{i} + 3\hat{j} - 4\hat{k}$ is :

- Options
1. 7
 2. $4\sqrt{3}$

3. 6

4. $2\sqrt{13}$

Question Type : MCQ

Question ID : 41652914400

Option 1 ID : 41652956379

Option 2 ID : 41652956381

Option 3 ID : 41652956378

Option 4 ID : 41652956380

Status : Answered

Chosen Option : 4

Q.13 The area (in sq. units) of the region bounded by the curves $y=2^x$ and $y=|x+1|$, in the first quadrant is :

Options

1. $\log_e 2 + \frac{3}{2}$

2. $\frac{3}{2}$

3. $\frac{1}{2}$

4. $\frac{3}{2} - \frac{1}{\log_e 2}$

Question Type : MCQ

Question ID : 41652914391

Option 1 ID : 41652956342

Option 2 ID : 41652956345

Option 3 ID : 41652956344

Option 4 ID : 41652956343

Status : Answered

Chosen Option : 4

Q.14 A spherical iron ball of radius 10 cm is coated with a layer of ice of uniform thickness that melts at a rate of $50 \text{ cm}^3/\text{min}$. When the thickness of the ice is 5 cm, then the rate at which the thickness (in cm/min) of the ice decreases, is :

Options

1. $\frac{1}{18\pi}$

2. $\frac{1}{36\pi}$

3. $\frac{5}{6\pi}$

4. $\frac{1}{9\pi}$

Question Type : MCQ

Question ID : 41652914387

Option 1 ID : **41652956328**
 Option 2 ID : **41652956326**
 Option 3 ID : **41652956327**
 Option 4 ID : **41652956329**
 Status : **Not Answered**
 Chosen Option : --

Q.15 If $\int x^5 e^{-x^2} dx = g(x)e^{-x^2} + c$, where c is a constant of integration, then $g(-1)$ is equal to :

- Options
1. -1
 2. 1
 3. $-\frac{5}{2}$
 4. $-\frac{1}{2}$

Question Type : **MCQ**
 Question ID : **41652914389**
 Option 1 ID : **41652956335**
 Option 2 ID : **41652956337**
 Option 3 ID : **41652956334**
 Option 4 ID : **41652956336**
 Status : **Not Attempted and Marked For Review**
 Chosen Option : --

Q.16 The smallest natural number n , such that the coefficient of x in the expansion of

$$\left(x^2 + \frac{1}{x^3}\right)^n \text{ is } {}^nC_{23}, \text{ is :}$$

- Options
1. 38
 2. 58
 3. 23
 4. 35

Question Type : **MCQ**
 Question ID : **41652914382**
 Option 1 ID : **41652956307**
 Option 2 ID : **41652956309**
 Option 3 ID : **41652956306**
 Option 4 ID : **41652956308**
 Status : **Answered**
 Chosen Option : **2**

Q.17 The sum of the real roots of the equation

$$\begin{vmatrix} x & -6 & -1 \\ 2 & -3x & x-3 \\ -3 & 2x & x+2 \end{vmatrix} = 0, \text{ is equal to :}$$

- Options
1. 6
 2. 0

3. 1
4. -4

Question Type : **MCQ**
 Question ID : **41652914379**
 Option 1 ID : **41652956297**
 Option 2 ID : **41652956295**
 Option 3 ID : **41652956296**
 Option 4 ID : **41652956294**
 Status : **Answered**
 Chosen Option : **3**

Q.18 Minimum number of times a fair coin must be tossed so that the probability of getting at least one head is more than 99% is :

- Options
1. 5
 2. 6
 3. 8
 4. 7

Question Type : **MCQ**
 Question ID : **41652914402**
 Option 1 ID : **41652956386**
 Option 2 ID : **41652956387**
 Option 3 ID : **41652956389**
 Option 4 ID : **41652956388**
 Status : **Answered**
 Chosen Option : **2**

Q.19 The angles A, B and C of a triangle ABC are in A.P. and $a : b = 1 : \sqrt{3}$. If $c = 4$ cm, then the area (in sq.cm) of this triangle is :

- Options
1. $\frac{2}{\sqrt{3}}$
 2. $4\sqrt{3}$
 3. $2\sqrt{3}$
 4. $\frac{4}{\sqrt{3}}$

Question Type : **MCQ**
 Question ID : **41652914404**
 Option 1 ID : **41652956397**
 Option 2 ID : **41652956394**
 Option 3 ID : **41652956395**
 Option 4 ID : **41652956396**
 Status : **Answered**
 Chosen Option : **3**

Q.20 The negation of the Boolean expression $\sim s \vee (\sim r \wedge s)$ is equivalent to :

- Options
1. $\sim s \wedge \sim r$

2. $\mathbf{r}$
3. $\mathbf{s} \cdot \mathbf{r}$
4. $\mathbf{s} \wedge \mathbf{r}$

Question Type : **MCQ**
 Question ID : **41652914405**
 Option 1 ID : **41652956401**
 Option 2 ID : **41652956400**
 Option 3 ID : **41652956398**
 Option 4 ID : **41652956399**
 Status : **Answered**
 Chosen Option : **4**

Q.21 A perpendicular is drawn from a point on the line $\frac{x-1}{2} = \frac{y+1}{-1} = \frac{z}{1}$ to the plane $x+y+z=3$ such that the foot of the perpendicular Q also lies on the plane $x-y+z=3$. Then the co-ordinates of Q are :

- Options**
1. (1, 0, 2)
 2. (2, 0, 1)
 3. (-1, 0, 4)
 4. (4, 0, -1)

Question Type : **MCQ**
 Question ID : **41652914398**
 Option 1 ID : **41652956371**
 Option 2 ID : **41652956370**
 Option 3 ID : **41652956372**
 Option 4 ID : **41652956373**
 Status : **Not Answered**
 Chosen Option : --

Q.22 Let a, b and c be in G.P. with common ratio r, where $a \neq 0$ and $0 < r \leq \frac{1}{2}$. If 3a, 7b and 15c are the first three terms of an A.P., then the 4th term of this A.P. is :

- Options**
1. $\frac{2}{3}a$
 2. 5a
 3. $\frac{7}{3}a$
 4. a

Question Type : **MCQ**
 Question ID : **41652914383**
 Option 1 ID : **41652956310**
 Option 2 ID : **41652956313**
 Option 3 ID : **41652956312**
 Option 4 ID : **41652956311**
 Status : **Answered**

Chosen Option : 3

Q.23 Lines are drawn parallel to the line $4x - 3y + 2 = 0$, at a distance $\frac{3}{5}$ from the origin. Then which one of the following points lies on any of these lines ?

Options

1. $\left(-\frac{1}{4}, \frac{2}{3}\right)$
2. $\left(\frac{1}{4}, -\frac{1}{3}\right)$
3. $\left(\frac{1}{4}, \frac{1}{3}\right)$
4. $\left(-\frac{1}{4}, -\frac{2}{3}\right)$

Question Type : MCQ

Question ID : 41652914393

Option 1 ID : 41652956353

Option 2 ID : 41652956351

Option 3 ID : 41652956350

Option 4 ID : 41652956352

Status : Answered

Chosen Option : 1

Q.24 If both the mean and the standard deviation of 50 observations $x_1, x_2, \dots, x_{50}$ are equal to 16, then the mean of $(x_1 - 4)^2, (x_2 - 4)^2, \dots, (x_{50} - 4)^2$ is :

Options

1. 400
2. 380
3. 525
4. 480

Question Type : MCQ

Question ID : 41652914401

Option 1 ID : 41652956383

Option 2 ID : 41652956382

Option 3 ID : 41652956385

Option 4 ID : 41652956384

Status : Answered

Chosen Option : 4

Q.25 If $\lim_{x \rightarrow 1} \frac{x^2 - ax + b}{x - 1} = 5$, then $a + b$ is equal to :

Options

1. -4
2. 5
3. -7

4. 1

Question Type : **MCQ**Question ID : **41652914385**Option 1 ID : **41652956320**Option 2 ID : **41652956319**Option 3 ID : **41652956321**Option 4 ID : **41652956318**Status : **Answered**Chosen Option : **3**

Q.26 Let $a_1, a_2, a_3, \dots$ be an A.P. with $a_6 = 2$. Then the common difference of this A.P., which maximises the product $a_1 a_4 a_5$, is :

- Options
1. $\frac{3}{2}$
 2. $\frac{8}{5}$
 3. $\frac{6}{5}$
 4. $\frac{2}{3}$

Question Type : **MCQ**Question ID : **41652914386**Option 1 ID : **41652956325**Option 2 ID : **41652956323**Option 3 ID : **41652956324**Option 4 ID : **41652956322**Status : **Not Answered**Chosen Option : **--**

Q.27 The integral $\int_{\pi/6}^{\pi/3} \sec^{2/3} x \operatorname{cosec}^{4/3} x \, dx$ is equal to :

- Options
1. $3^{5/6} - 3^{2/3}$
 2. $3^{4/3} - 3^{1/3}$
 3. $3^{7/6} - 3^{5/6}$
 4. $3^{5/3} - 3^{1/3}$

Question Type : **MCQ**Question ID : **41652914390**Option 1 ID : **41652956340**Option 2 ID : **41652956339**Option 3 ID : **41652956341**Option 4 ID : **41652956338**Status : **Not Answered**Chosen Option : **--****Q.28**

If the plane $2x - y + 2z + 3 = 0$ has the distances $\frac{1}{3}$ and $\frac{2}{3}$ units from the planes $4x - 2y + 4z + \lambda = 0$ and $2x - y + 2z + \mu = 0$, respectively, then the maximum value of $\lambda + \mu$ is equal to :

- Options
1. 9
 2. 15
 3. 5
 4. 13

Question Type : **MCQ**
 Question ID : **41652914399**
 Option 1 ID : **41652956375**
 Option 2 ID : **41652956377**
 Option 3 ID : **41652956374**
 Option 4 ID : **41652956376**
 Status : **Answered**
 Chosen Option : **3**

Q.29 Let $y = y(x)$ be the solution of the differential equation, $\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, such that $y(0) = 1$. Then :

- Options
1. $y\left(\frac{\pi}{4}\right) + y\left(-\frac{\pi}{4}\right) = \frac{\pi^2}{2} + 2$
 2. $y'\left(\frac{\pi}{4}\right) + y'\left(-\frac{\pi}{4}\right) = -\sqrt{2}$
 3. $y\left(\frac{\pi}{4}\right) - y\left(-\frac{\pi}{4}\right) = \sqrt{2}$
 4. $y'\left(\frac{\pi}{4}\right) - y'\left(-\frac{\pi}{4}\right) = \pi - \sqrt{2}$

Question Type : **MCQ**
 Question ID : **41652914392**
 Option 1 ID : **41652956347**
 Option 2 ID : **41652956349**
 Option 3 ID : **41652956346**
 Option 4 ID : **41652956348**
 Status : **Answered**
 Chosen Option : **3**

Q.30 The locus of the centres of the circles, which touch the circle, $x^2 + y^2 = 1$ externally, also touch the y -axis and lie in the first quadrant, is :

- Options
1. $x = \sqrt{1+4y}, y \geq 0$

2. $y = \sqrt{1+2x}, x \geq 0$

3. $y = \sqrt{1+4x}, x \geq 0$

4. $x = \sqrt{1+2y}, y \geq 0$

Question Type : **MCQ**Question ID : **41652914394**Option 1 ID : **41652956356**Option 2 ID : **41652956354**Option 3 ID : **41652956355**Option 4 ID : **41652956357**Status : **Answered**Chosen Option : **3**