

JEE April 2019

Roll No.	
Candidate Name	
Application No	
Test Date	12/04/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

Q.1 Two particles are projected from the same point with the same speed u such that they have the same range R , but different maximum heights, h_1 and h_2 . Which of the following is correct ?

- Options**
1. $R^2 = 4 h_1 h_2$
 2. $R^2 = 16 h_1 h_2$
 3. $R^2 = 2 h_1 h_2$
 4. $R^2 = h_1 h_2$

Question Type : **MCQ**

Question ID : **41652913418**

Option 1 ID : **41652952451**

Option 2 ID : **41652952450**

Option 3 ID : **41652952452**

Option 4 ID : **41652952453**

Status : **Not Answered**

Chosen Option : --

Q.2 In an amplitude modulator circuit, the carrier wave is given by,
 $C(t) = 4 \sin(20000 \pi t)$ while modulating signal is given by, $m(t) = 2 \sin(2000 \pi t)$. The values of modulation index and lower side band frequency are :

- Options**
1. 0.5 and 10 kHz
 2. 0.4 and 10 kHz
 3. 0.3 and 9 kHz
 4. 0.5 and 9 kHz

Question Type : **MCQ**

Question ID : **41652913443**

Option 1 ID : **41652952552**

Option 2 ID : **41652952551**

Option 3 ID : **41652952550**

Option 4 ID : **41652952553**

Status : **Answered**

Chosen Option : **1**

Q.3 Two sources of sound S_1 and S_2 produce sound waves of same frequency 660 Hz. A listener is moving from source S_1 towards S_2 with a constant speed u m/s and he hears 10 beats/s. The velocity of sound is 330 m/s. Then, u equals :

- Options
1. 5.5 m/s
 2. 15.0 m/s
 3. 2.5 m/s
 4. 10.0 m/s

Question Type : **MCQ**

Question ID : **41652913429**

Option 1 ID : **41652952495**

Option 2 ID : **41652952497**


Option 3 ID : **41652952494**

Option 4 ID : **41652952496**

Status : **Not Answered**

Chosen Option : --

Q.4 Three particles of masses 50 g, 100 g and 150 g are placed at the vertices of an equilateral triangle of side 1 m (as shown in the figure). The (x, y) coordinates of the centre of mass will be :



- Options
1. $\left(\frac{\sqrt{3}}{4} \text{ m}, \frac{5}{12} \text{ m}\right)$
 2. $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{8} \text{ m}\right)$
 3. $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{4} \text{ m}\right)$
 4. $\left(\frac{\sqrt{3}}{8} \text{ m}, \frac{7}{12} \text{ m}\right)$

Question Type : **MCQ**

Question ID : **41652913422**

Option 1 ID : **41652952468**

Option 2 ID : **41652952466**

Option 3 ID : **41652952467**

Option 4 ID : **41652952469**
 Status : **Marked For Review**
 Chosen Option : **3**

Q.5 A Carnot engine has an efficiency of $1/6$. When the temperature of the sink is reduced by 62°C , its efficiency is doubled. The temperatures of the source and the sink are, respectively,

- Options
1. 62°C , 124°C
 2. 99°C , 37°C
 3. 124°C , 62°C
 4. 37°C , 99°C

Question Type : **MCQ**
 Question ID : **41652913426**
 Option 1 ID : **41652952485**
 Option 2 ID : **41652952482**
 Option 3 ID : **41652952483**
 Option 4 ID : **41652952484**
 Status : **Answered**
 Chosen Option : **2**


Q.6 A spring whose unstretched length is l has a force constant k . The spring is cut into two pieces of unstretched lengths l_1 and l_2 where, $l_1 = nl_2$ and n is an integer. The ratio k_1/k_2 of the corresponding force constants, k_1 and k_2 will be :

- Options
1. n
 2. $\frac{1}{n^2}$
 3. $\frac{1}{n}$
 4. n^2

Question Type : **MCQ**
 Question ID : **41652913420**
 Option 1 ID : **41652952458**
 Option 2 ID : **41652952461**
 Option 3 ID : **41652952459**
 Option 4 ID : **41652952460**
 Status : **Answered**
 Chosen Option : **4**

Q.7

A transparent cube of side d , made of a material of refractive index μ_2 , is immersed in a liquid of refractive index μ_1 ($\mu_1 < \mu_2$). A ray is incident on the face AB at an angle θ (shown in the figure). Total internal reflection takes place at point E on the face BC.



Then θ must satisfy :

Options

1. $\theta < \sin^{-1} \frac{\mu_1}{\mu_2}$
2. $\theta > \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$
3. $\theta < \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$
4. $\theta > \sin^{-1} \frac{\mu_1}{\mu_2}$

Question Type : **MCQ**

Question ID : **41652913437**

Option 1 ID : **41652952529**

Option 2 ID : **41652952527**

Option 3 ID : **41652952526**

Option 4 ID : **41652952528**

Status : **Answered**

Chosen Option : **2**

Q.8 A tuning fork of frequency 480 Hz is used in an experiment for measuring speed of sound (v) in air by resonance tube method. Resonance is observed to occur at two successive lengths of the air column, $l_1 = 30$ cm and $l_2 = 70$ cm. Then, v is equal to :

Options

1. 332 ms^{-1}
2. 384 ms^{-1}
3. 338 ms^{-1}
4. 379 ms^{-1}

Question Type : **MCQ**

Question ID : 41652913444

Option 1 ID : 41652952557

Option 2 ID : 41652952556

Option 3 ID : 41652952554

Option 4 ID : 41652952555

Status : Not Answered

Chosen Option : --

Q.9 The electron in a hydrogen atom first jumps from the third excited state to the second excited state and subsequently to the first excited state. The ratio of the respective wavelengths, λ_1/λ_2 , of the photons emitted in this process is :

- Options
1. 20/7
 2. 27/5
 3. 7/5
 4. 9/7

Question Type : MCQ

Question ID : 41652913440

Option 1 ID : 41652952541

Option 2 ID : 41652952538

Option 3 ID : 41652952539

Option 4 ID : 41652952540

Status : Answered

Chosen Option : 2

Q.10 A diatomic gas with rigid molecules does 10 J of work when expanded at constant pressure. What would be the heat energy absorbed by the gas, in this process ?

- Options
1. 25 J
 2. 35 J
 3. 30 J
 4. 40 J

Question Type : MCQ

Question ID : 41652913427

Option 1 ID : 41652952486

Option 2 ID : 41652952488

Option 3 ID : 41652952487

Option 4 ID : 41652952489

Status : Marked For Review

Chosen Option : 1

Q.11

Let a total charge $2Q$ be distributed in a sphere of radius R , with the charge density given by $\rho(r) = kr$, where r is the distance from the centre. Two charges A and B , of $-Q$ each, are placed on diametrically opposite points, at equal distance, a , from the centre. If A and B do not experience any force, then :

- Options
1. $a = 8^{-1/4} R$
 2. $a = \frac{3R}{2^{1/4}}$
 3. $a = 2^{-1/4} R$
 4. $a = R/\sqrt{3}$

Question Type : **MCQ**

Question ID : **41652913430**

Option 1 ID : **41652952498**

Option 2 ID : **41652952501**

Option 3 ID : **41652952499**

Option 4 ID : **41652952500**

Status : **Not Answered**

Chosen Option : --

Q.12 Consider an electron in a hydrogen atom, revolving in its second excited state (having radius $4.65 \text{ \AA}$). The de-Broglie wavelength of this electron is :

- Options
1. $3.5 \text{ \AA}$
 2. $6.6 \text{ \AA}$
 3. $12.9 \text{ \AA}$
 4. $9.7 \text{ \AA}$

Question Type : **MCQ**

Question ID : **41652913439**

Option 1 ID : **41652952534**

Option 2 ID : **41652952535**

Option 3 ID : **41652952537**

Option 4 ID : **41652952536**

Status : **Not Attempted and Marked For Review**

Chosen Option : --

Q.13 A solid sphere, of radius R acquires a terminal velocity v_1 when falling (due to gravity) through a viscous fluid having a coefficient of viscosity η . The sphere is broken into 27 identical solid spheres. If each of these spheres acquires a terminal velocity, v_2 , when falling through the same fluid, the ratio (v_1/v_2) equals :

Options 1. 9

2. $1/27$ 3. $1/9$

4. 27

Question Type : MCQ

Question ID : 41652913425

Option 1 ID : 41652952478

Option 2 ID : 41652952481


Option 3 ID : 41652952479

Option 4 ID : 41652952480

Status : Not Answered

Chosen Option : --

Q.14 A smooth wire of length $2\pi r$ is bent into a circle and kept in a vertical plane. A bead can slide smoothly on the wire. When the circle is rotating with angular speed ω about the vertical diameter AB, as shown in figure, the bead is at rest with respect to the circular ring at position P as shown. Then the value of ω^2 is equal to :

Options 1. $\frac{\sqrt{3}g}{2r}$ 2. $2g/(r\sqrt{3})$ 3. $(g\sqrt{3})/r$ 4. $2g/r$

Question Type : MCQ

Question ID : 41652913421

Option 1 ID : 41652952463

Option 2 ID : 41652952465

Option 3 ID : 41652952464

Option 4 ID : 41652952462

Status : Not Answered

Chosen Option : --

Q.15

A particle is moving with speed $v = b\sqrt{x}$ along positive x -axis. Calculate the speed of the particle at time $t = \tau$ (assume that the particle is at origin at $t = 0$).

Options

1. $\frac{b^2\tau}{4}$
2. $\frac{b^2\tau}{2}$
3. $b^2\tau$
4. $\frac{b^2\tau}{\sqrt{2}}$

Question Type : MCQ

Question ID : 41652913417

Option 1 ID : 41652952446

Option 2 ID : 41652952448

Option 3 ID : 41652952449

Option 4 ID : 41652952447

Status : Answered

Chosen Option : 1

Q.16

The ratio of the weights of a body on the Earth's surface to that on the surface of a planet is 9 : 4. The mass of the planet is $\frac{1}{9}$ th of that of the Earth. If 'R' is the radius of the Earth, what is the radius of the planet ? (Take the planets to have the same mass density)

Options

1. $\frac{R}{3}$
2. $\frac{R}{4}$
3. $\frac{R}{9}$
4. $\frac{R}{2}$

Question Type : MCQ

Question ID : 41652913423

Option 1 ID : 41652952471

Option 2 ID : 41652952473

Option 3 ID : 41652952472

Option 4 ID : 41652952470

Status : Answered

Chosen Option : 3

Q.17

A system of three polarizers P_1, P_2, P_3 is set up such that the pass axis of P_3 is crossed with respect to that of P_1 . The pass axis of P_2 is inclined at 60° to the pass axis of P_3 . When a beam of unpolarized light of intensity I_0 is incident on P_1 , the intensity of light transmitted by the three polarizers is I . The ratio (I_0/I) equals (nearly) :

- Options
1. 5.33
 2. 16.00
 3. 10.67
 4. 1.80

Question Type : **MCQ**

Question ID : **41652913438**

Option 1 ID : **41652952531**

Option 2 ID : **41652952533**

Option 3 ID : **41652952532**

Option 4 ID : **41652952530**

Status : **Answered**

Chosen Option : **3**

Q.18 A uniform cylindrical rod of length L and radius r , is made from a material whose Young's modulus of Elasticity equals Y . When this rod is heated by temperature T and simultaneously subjected to a net longitudinal compressional force F , its length remains unchanged. The coefficient of volume expansion, of the material of the rod, is (nearly) equal to :

- Options
1. $9F/(\pi r^2 Y T)$
 2. $6F/(\pi r^2 Y T)$
 3. $3F/(\pi r^2 Y T)$
 4. $F/(3\pi r^2 Y T)$

Question Type : **MCQ**

Question ID : **41652913424**

Option 1 ID : **41652952475**

Option 2 ID : **41652952474**

Option 3 ID : **41652952476**

Option 4 ID : **41652952477**

Status : **Not Answered**

Chosen Option : **--**

Q.19

The number density of molecules of a gas depends on their distance r from the origin as, $n(r) = n_0 e^{-\alpha r^4}$. Then the total number of molecules is proportional to :

Options

1. $n_0 \alpha^{-3/4}$
2. $\sqrt{n_0} \alpha^{1/2}$
3. $n_0 \alpha^{1/4}$
4. $n_0 \alpha^{-3}$

Question Type : MCQ

Question ID : 41652913416

Option 1 ID : 41652952444

Option 2 ID : 41652952443

Option 3 ID : 41652952445

Option 4 ID : 41652952442

Status : Not Attempted and Marked For Review

Chosen Option : --

Q.20 A small speaker delivers 2 W of audio output. At what distance from the speaker will one detect 120 dB intensity sound ? [Given reference intensity of sound as 10^{-12} W/m^2]

Options

1. 40 cm
2. 20 cm
3. 10 cm
4. 30 cm

Question Type : MCQ

Question ID : 41652913428

Option 1 ID : 41652952493

Option 2 ID : 41652952491

Option 3 ID : 41652952490

Option 4 ID : 41652952492

Status : Not Answered

Chosen Option : --

Q.21 Half lives of two radioactive nuclei A and B are 10 minutes and 20 minutes, respectively. If, initially a sample has equal number of nuclei, then after 60 minutes, the ratio of decayed numbers of nuclei A and B will be :

Options

1. 3 : 8
2. 1 : 8
3. 8 : 1
4. 9 : 8

Question Type : MCQ

Question ID : 41652913441

Option 1 ID : 41652952545

Option 2 ID : 41652952542


Option 3 ID : 41652952544

Option 4 ID : 41652952543

Status : Answered

Chosen Option : 2

- Q.22** An electron, moving along the x -axis with an initial energy of 100 eV, enters a region of magnetic field $\vec{B} = (1.5 \times 10^{-3} \text{ T}) \hat{k}$ at S (See figure). The field extends between $x=0$ and $x=2$ cm. The electron is detected at the point Q on a screen placed 8 cm away from the point S. The distance d between P and Q (on the screen) is :
(electron's charge = $1.6 \times 10^{-19} \text{ C}$, mass of electron = $9.1 \times 10^{-31} \text{ kg}$)



- Options
1. 11.65 cm
 2. 12.87 cm
 3. 1.22 cm
 4. 2.25 cm

Question Type : MCQ

Question ID : 41652913433

Option 1 ID : 41652952512

Option 2 ID : 41652952511

Option 3 ID : 41652952510


Option 4 ID : 41652952513

Status : Not Attempted and Marked For Review

Chosen Option : --

Q.23

In the given circuit, the charge on $4\ \mu\text{F}$ capacitor will be :



- Options
1. $5.4\ \mu\text{C}$
 2. $9.6\ \mu\text{C}$
 3. $13.4\ \mu\text{C}$
 4. $24\ \mu\text{C}$

Question Type : **MCQ**

Question ID : **41652913431**

Option 1 ID : **41652952504**

Option 2 ID : **41652952502**

Option 3 ID : **41652952505**

Option 4 ID : **41652952503**

Status : **Answered**

Chosen Option : **1**

Q.24 One kg of water, at 20°C , is heated in an electric kettle whose heating element has a mean (temperature averaged) resistance of $20\ \Omega$. The rms voltage in the mains is $200\ \text{V}$. Ignoring heat loss from the kettle, time taken for water to evaporate fully, is close to :

[Specific heat of water = $4200\ \text{J}/(\text{kg } ^\circ\text{C})$,
Latent heat of water = $2260\ \text{kJ}/\text{kg}$]

- Options
1. 16 minutes
 2. 22 minutes
 3. 3 minutes
 4. 10 minutes

Question Type : **MCQ**

Question ID : **41652913432**

Option 1 ID : **41652952509**

Option 2 ID : **41652952508**

Option 3 ID : **41652952506**

Option 4 ID : **41652952507**

Status : **Answered**

Chosen Option : **3**

Q.25

A moving coil galvanometer, having a resistance G , produces full scale deflection when a current I_g flows through it. This galvanometer can be converted into
 (i) an ammeter of range 0 to I_0 ($I_0 > I_g$) by connecting a shunt resistance R_A to it and
 (ii) into a voltmeter of range 0 to V ($V = GI_0$) by connecting a series resistance R_V to it.
 Then,

Options

$$R_A R_V = G^2 \left(\frac{I_0 - I_g}{I_g} \right) \text{ and}$$

1.

$$\frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$$

$$2. R_A R_V = G^2 \text{ and } \frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$$

$$R_A R_V = G^2 \left(\frac{I_g}{I_0 - I_g} \right) \text{ and}$$

3.

$$\frac{R_A}{R_V} = \left(\frac{I_0 - I_g}{I_g} \right)^2$$

$$4. R_A R_V = G^2 \text{ and } \frac{R_A}{R_V} = \frac{I_g}{I_0 - I_g}$$

Question Type : MCQ

Question ID : 41652913445

Option 1 ID : 41652952561

Option 2 ID : 41652952558

Option 3 ID : 41652952560

Option 4 ID : 41652952559


Status : Answered

Chosen Option : 3

Q.26

Find the magnetic field at point P due to a straight line segment AB of length 6 cm carrying a current of 5 A. (See figure)

$$(\mu_0 = 4\pi \times 10^{-7} \text{ N-A}^{-2})$$



- Options
1. $2.0 \times 10^{-5} \text{ T}$
 2. $1.5 \times 10^{-5} \text{ T}$
 3. $3.0 \times 10^{-5} \text{ T}$
 4. $2.5 \times 10^{-5} \text{ T}$

Question Type : MCQ

Question ID : 41652913434

Option 1 ID : 41652952515

Option 2 ID : 41652952514

Option 3 ID : 41652952517

Option 4 ID : 41652952516

Status : Answered

Chosen Option : 1

Q.27 A plane electromagnetic wave having a frequency $\nu = 23.9 \text{ GHz}$ propagates along the positive z-direction in free space. The peak value of the Electric Field is 60 V/m. Which among the following is the acceptable magnetic field component in the electromagnetic wave ?

- Options
1. $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z + 1.5 \times 10^{11} t) \hat{i}$
 2. $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z - 1.5 \times 10^{11} t) \hat{i}$
 3. $\vec{B} = 60 \sin(0.5 \times 10^3 x + 1.5 \times 10^{11} t) \hat{k}$
 4. $\vec{B} = 2 \times 10^{-7} \sin(1.5 \times 10^2 x + 0.5 \times 10^{11} t) \hat{j}$

Question Type : MCQ

Question ID : 41652913436

Option 1 ID : 41652952524

Option 2 ID : 41652952523


Option 3 ID : 41652952522

Option 4 ID : 41652952525

Status : Not Answered

Chosen Option : --

- Q.28** Consider the LR circuit shown in the figure.
If the switch S is closed at $t=0$ then the amount of charge that passes through the battery between $t=0$ and $t=\frac{L}{R}$ is :



Options

1. $\frac{2.7 EL}{R^2}$
2. $\frac{EL}{2.7R^2}$
3. $\frac{7.3 EL}{R^2}$
4. $\frac{EL}{7.3R^2}$

Question Type : MCQ

Question ID : 41652913435

Option 1 ID : 41652952519

Option 2 ID : 41652952520


Option 3 ID : 41652952518

Option 4 ID : 41652952521

Status : Not Answered

Chosen Option : --

- Q.29** Figure shows a DC voltage regulator circuit, with a Zener diode of breakdown voltage = 6V. If the unregulated input voltage varies between 10 V to 16 V, then what is the maximum Zener current ?



Options

1. 2.5 mA
2. 1.5 mA

3. 7.5 mA
4. 3.5 mA

Question Type : MCQ

Question ID : 41652913442

Option 1 ID : 41652952548

Option 2 ID : 41652952547


Option 3 ID : 41652952546

Option 4 ID : 41652952549

Status : Not Answered

Chosen Option : --

Q.30 A block of mass 5 kg is (i) pushed in case (A) and (ii) pulled in case (B), by a force $F = 20$ N, making an angle of 30° with the horizontal, as shown in the figures. The coefficient of friction between the block and floor is $\mu = 0.2$. The difference between the accelerations of the block, in case (B) and case (A) will be : ($g = 10 \text{ ms}^{-2}$)



- Options
1. 0.4 ms^{-2}
 2. 3.2 ms^{-2}
 3. 0.8 ms^{-2}
 4. 0 ms^{-2}

Question Type : MCQ

Question ID : 41652913419

Option 1 ID : 41652952456

Option 2 ID : 41652952455

Option 3 ID : 41652952457

Option 4 ID : 41652952454

Status : Not Answered

Chosen Option : --

Section : Chemistry


Q.1 Thermal decomposition of a Mn compound (X) at 513 K results in compound Y, MnO_2 and a gaseous product. MnO_2 reacts with NaCl and concentrated H_2SO_4 to give a pungent gas Z. X, Y, and Z, respectively, are :

- Options
1. K_3MnO_4 , K_2MnO_4 and Cl_2
 2. K_2MnO_4 , KMnO_4 and SO_2

3. KMnO_4 , K_2MnO_4 and Cl_2
4. K_2MnO_4 , KMnO_4 and Cl_2

Question Type : **MCQ**Question ID : **41652913461**Option 1 ID : **41652952625**Option 2 ID : **41652952622**Option 3 ID : **41652952623**Option 4 ID : **41652952624**Status : **Answered**Chosen Option : **2**

Q.2 NO_2 required for a reaction is produced by the decomposition of N_2O_5 in CCl_4 as per the equation,



The initial concentration of N_2O_5 is 3.00 mol L^{-1} and it is 2.75 mol L^{-1} after 30 minutes. The rate of formation of NO_2 is :

- Options
1. $4.167 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
 2. $1.667 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$
 3. $8.333 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$
 4. $2.083 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

Question Type : **MCQ**Question ID : **41652913474**Option 1 ID : **41652952674**Option 2 ID : **41652952677**Option 3 ID : **41652952675**Option 4 ID : **41652952676**Status : **Not Answered**Chosen Option : **--**

Q.3 The pair that has similar atomic radii is :

- Options
1. Mn and Re
 2. Ti and Hf
 3. Sc and Ni
 4. Mo and W

Question Type : **MCQ**Question ID : **41652913462**Option 1 ID : **41652952627**Option 2 ID : **41652952626**Option 3 ID : **41652952628**Option 4 ID : **41652952629**Status : **Answered**Chosen Option : **4**

Q.4 The INCORRECT statement is :

Options

- Lithium is the strongest reducing agent among the alkali metals.
- Lithium is least reactive with water among the alkali metals.
- LiNO_3 decomposes on heating to give LiNO_2 and O_2 .
- LiCl crystallises from aqueous solution as $\text{LiCl} \cdot 2\text{H}_2\text{O}$.

Question Type : MCQ

Question ID : 41652913459

Option 1 ID : 41652952615

Option 2 ID : 41652952616

Option 3 ID : 41652952614

Option 4 ID : 41652952617

Status : Answered

Chosen Option : 4

Q.5 The C – C bond length is maximum in :

- Options
1. graphite
 2. C_{70}
 3. C_{60}
 4. diamond

Question Type : MCQ

Question ID : 41652913460

Option 1 ID : 41652952619

Option 2 ID : 41652952621

Option 3 ID : 41652952618

Option 4 ID : 41652952620

Status : Answered

Chosen Option : 4

Q.6 A solution is prepared by dissolving 0.6 g of urea (molar mass = 60 g mol^{-1}) and 1.8 g of glucose (molar mass = 180 g mol^{-1}) in 100 mL of water at 27°C . The osmotic pressure of the solution is :
($R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$)

- Options
1. 8.2 atm
 2. 2.46 atm
 3. 4.92 atm
 4. 1.64 atm

Question Type : MCQ

Question ID : 41652913470

Option 1 ID : 41652952661

Option 2 ID : 41652952659

Option 3 ID : 41652952658

Option 4 ID : **41652952660**Status : **Answered**Chosen Option : **3**

Q.7 In comparison to boron, beryllium has :

Options

1. lesser nuclear charge and lesser first ionisation enthalpy.
2. greater nuclear charge and lesser first ionisation enthalpy.
3. greater nuclear charge and greater first ionisation enthalpy.
4. lesser nuclear charge and greater first ionisation enthalpy.

Question Type : **MCQ**Question ID : **41652913456**Option 1 ID : **41652952604**Option 2 ID : **41652952605**Option 3 ID : **41652952602**Option 4 ID : **41652952603**Status : **Answered**Chosen Option : **4**

Q.8 The decreasing order of electrical conductivity of the following aqueous solutions is :

0.1 M Formic acid (A),

0.1 M Acetic acid (B),

0.1 M Benzoic acid (C).


Options

1. $A > C > B$
2. $C > B > A$
3. $A > B > C$
4. $C > A > B$

Question Type : **MCQ**Question ID : **41652913473**Option 1 ID : **41652952673**Option 2 ID : **41652952672**Option 3 ID : **41652952670**Option 4 ID : **41652952671**Status : **Answered**Chosen Option : **3**

Q.9 What will be the major product when m-cresol is reacted with propargyl bromide ($\text{HC} \equiv \text{C}-\text{CH}_2\text{Br}$) in presence of K_2CO_3 in acetone ?

Options

Question Type : **MCQ**Question ID : **41652913453**Option 1 ID : **41652952590**Option 2 ID : **41652952591**Option 3 ID : **41652952592**Option 4 ID : **41652952593**Status : **Not Answered**

Chosen Option : --

Q.10 The molar solubility of $\text{Cd}(\text{OH})_2$ is $1.84 \times 10^{-5} \text{ M}$ in water. The expected solubility of $\text{Cd}(\text{OH})_2$ in a buffer solution of $\text{pH} = 12$ is :

Options

1. $1.84 \times 10^{-9} \text{ M}$

2. $\frac{2.49}{1.84} \times 10^{-9} \text{ M}$

3. $6.23 \times 10^{-11} \text{ M}$

4. $2.49 \times 10^{-10} \text{ M}$





Question Type : **MCQ**Question ID : **41652913472**Option 1 ID : **41652952669**Option 2 ID : **41652952668**Option 3 ID : **41652952666**Option 4 ID : **41652952667**Status : **Not Answered**

Chosen Option : --

Q.11

Benzene diazonium chloride on reaction with aniline in the presence of dilute hydrochloric acid gives :

Options

1. 
2. 
3. 
4. 

Question Type : **MCQ**

Question ID : **41652913451**

Option 1 ID : **41652952585**

Option 2 ID : **41652952584**

Option 3 ID : **41652952582**

Option 4 ID : **41652952583**

Status : **Answered**

Chosen Option : **3**

Q.12 The correct statement is :

Options

1. leaching of bauxite using concentrated NaOH solution gives sodium aluminate and sodium silicate.
2. the Hall-Heroult process is used for the production of aluminium and iron.
3. pig iron is obtained from cast iron.
4. the blistered appearance of copper during the metallurgical process is due to the evolution of CO₂.

Question Type : **MCQ**

Question ID : **41652913457**

Option 1 ID : **41652952608**

Option 2 ID : **41652952609**

Option 3 ID : **41652952607**

Option 4 ID : **41652952606**

Status : **Answered**

Chosen Option : **3**

Q.13 The primary pollutant that leads to photochemical smog is :

Options

1. acrolein

2. nitrogen oxides
3. ozone
4. sulphur dioxide

Question Type : MCQ

Question ID : 41652913465

Option 1 ID : 41652952640

Option 2 ID : 41652952641


Option 3 ID : 41652952638

Option 4 ID : 41652952639

Status : Answered

Chosen Option : 2

Q.14 The IUPAC name for the following compound is :



- Options
1. 3-methyl-4-(3-methylprop-1-enyl)-1-heptyne
 2. 3,5-dimethyl-4-propylhept-6-en-1-yne
 3. 3-methyl-4-(1-methylprop-2-ynyl)-1-heptene
 4. 3,5-dimethyl-4-propylhept-1-en-6-yne

Question Type : MCQ

Question ID : 41652913447

Option 1 ID : 41652952567

Option 2 ID : 41652952566

Option 3 ID : 41652952569

Option 4 ID : 41652952568

Status : Answered

Chosen Option : 4

Q.15 The temporary hardness of a water sample is due to compound X. Boiling this sample converts X to compound Y. X and Y, respectively, are :

- Options
1. $\text{Mg}(\text{HCO}_3)_2$ and $\text{Mg}(\text{OH})_2$
 2. $\text{Ca}(\text{HCO}_3)_2$ and $\text{Ca}(\text{OH})_2$
 3. $\text{Mg}(\text{HCO}_3)_2$ and MgCO_3
 4. $\text{Ca}(\text{HCO}_3)_2$ and CaO

Question Type : MCQ

Question ID : 41652913458

Option 1 ID : 41652952610

Option 2 ID : 41652952612


Option 3 ID : 41652952611

Option 4 ID : 41652952613

Status : Answered

Chosen Option : 2

Q.16 In the following skew conformation of ethane, $H' - C - C - H''$ dihedral angle is :



- Options
1. 58°
 2. 149°
 3. 151°
 4. 120°

Question Type : MCQ

Question ID : 41652913446

Option 1 ID : 41652952562

Option 2 ID : 41652952564

Option 3 ID : 41652952563

Option 4 ID : 41652952565

Status : Not Answered

Chosen Option : --

Q.17 Among the following, the energy of 2s orbital is lowest in :

- Options
1. K
 2. H
 3. Li
 4. Na

Question Type : MCQ

Question ID : 41652913468

Option 1 ID : 41652952651

Option 2 ID : 41652952650

Option 3 ID : 41652952653

Option 4 ID : 41652952652

Status : Answered

Chosen Option : 3

Q.18 Which one of the following is likely to give a precipitate with $AgNO_3$ solution ?

Options

1. $\text{CH}_2=\text{CH}-\text{Cl}$
2. CCl_4
3. CHCl_3
4. $(\text{CH}_3)_3\text{CCl}$

Question Type : **MCQ**Question ID : **41652913455**Option 1 ID : **41652952601**Option 2 ID : **41652952598**Option 3 ID : **41652952599**Option 4 ID : **41652952600**Status : **Answered**Chosen Option : **1**




Q.19 25 g of an unknown hydrocarbon upon burning produces 88 g of CO_2 and 9 g of H_2O . This unknown hydrocarbon contains :

- Options
1. 20 g of carbon and 5 g of hydrogen
 2. 22 g of carbon and 3 g of hydrogen
 3. 24 g of carbon and 1 g of hydrogen
 4. 18 g of carbon and 7 g of hydrogen

Question Type : **MCQ**Question ID : **41652913466**Option 1 ID : **41652952642**Option 2 ID : **41652952644**Option 3 ID : **41652952643**Option 4 ID : **41652952645**Status : **Answered**Chosen Option : **3**

Q.20 Heating of 2-chloro-1-phenylbutane with EtOK/EtOH gives X as the major product. Reaction of X with $\text{Hg}(\text{OAc})_2/\text{H}_2\text{O}$ followed by NaBH_4 gives Y as the major product. Y is :

Options

1. 
2. 
3. 
4. 

Question Type : **MCQ**

Question ID : 41652913452

Option 1 ID : 41652952586

Option 2 ID : 41652952588

Option 3 ID : 41652952587

Option 4 ID : 41652952589

Status : Answered

Chosen Option : 1

Q.21 The compound used in the treatment of lead poisoning is :

- Options
1. D-penicillamine
 2. desferrioxime B
 3. Cis-platin
 4. EDTA

Question Type : MCQ

Question ID : 41652913463

Option 1 ID : 41652952630

Option 2 ID : 41652952633

Option 3 ID : 41652952632

Option 4 ID : 41652952631

Status : Answered

Chosen Option : 3

Q.22 Which of the given statements is INCORRECT about glycogen ?

- Options
1. It is a straight chain polymer similar to amylose.
 2. Only α -linkages are present in the molecule.
 3. It is present in animal cells.
 4. It is present in some yeast and fungi.

Question Type : MCQ

Question ID : 41652913449

Option 1 ID : 41652952575

Option 2 ID : 41652952577


Option 3 ID : 41652952574

Option 4 ID : 41652952576

Status : Answered

Chosen Option : 2

Q.23 Consider the following reactions :



'A' is :

- Options
1. $\text{CH} \equiv \text{CH}$

2. $\text{CH}_3-\text{C}\equiv\text{C}-\text{CH}_3$
3. $\text{CH}_3-\text{C}\equiv\text{CH}$
4. $\text{CH}_2=\text{CH}_2$

Question Type : MCQ

Question ID : 41652913448

Option 1 ID : 41652952570

Option 2 ID : 41652952572


Option 3 ID : 41652952571

Option 4 ID : 41652952573

Status : Answered

Chosen Option : 2

Q.24 The correct name of the following polymer is :



- Options
1. Polyisobutane
 2. Polytert-butylene
 3. Polyisoprene
 4. Polyisobutylene

Question Type : MCQ

Question ID : 41652913450

Option 1 ID : 41652952578

Option 2 ID : 41652952580

Option 3 ID : 41652952581

Option 4 ID : 41652952579

Status : Answered

Chosen Option : 4

Q.25 An 'Assertion' and a 'Reason' are given below. Choose the correct answer from the following options :

Assertion (A): Vinyl halides do not undergo nucleophilic substitution easily.

Reason (R): Even though the intermediate carbocation is stabilized by loosely held π -electrons, the cleavage is difficult because of strong bonding.

- Options
1. Both (A) and (R) are wrong statements.
 2. Both (A) and (R) are correct statements and (R) is the correct explanation of (A).

Both (A) and (R) are correct

3. statements but (R) is not the correct explanation of (A).
4. (A) is a correct statement but (R) is a wrong statement.

Question Type : MCQ

Question ID : 41652913454

Option 1 ID : 41652952597

Option 2 ID : 41652952594

Option 3 ID : 41652952595

Option 4 ID : 41652952596

Status : Answered

Chosen Option : 3

Q.26 The ratio of number of atoms present in a simple cubic, body centered cubic and face centered cubic structure are, respectively :

Options

1. 8 : 1 : 6
2. 1 : 2 : 4
3. 4 : 2 : 1
4. 4 : 2 : 3

Question Type : MCQ

Question ID : 41652913467

Option 1 ID : 41652952646

Option 2 ID : 41652952647

Option 3 ID : 41652952649

Option 4 ID : 41652952648

Status : Answered

Chosen Option : 2

Q.27 In which one of the following equilibria,
 $K_p \neq K_c$?

Options

1. $2\text{C(s)} + \text{O}_2\text{(g)} \rightleftharpoons 2\text{CO(g)}$
2. $2\text{HI(g)} \rightleftharpoons \text{H}_2\text{(g)} + \text{I}_2\text{(g)}$
3. $\text{NO}_2\text{(g)} + \text{SO}_2\text{(g)} \rightleftharpoons \text{NO(g)} + \text{SO}_3\text{(g)}$
4. $2\text{NO(g)} \rightleftharpoons \text{N}_2\text{(g)} + \text{O}_2\text{(g)}$

Question Type : MCQ

Question ID : 41652913471

Option 1 ID : 41652952664

Option 2 ID : 41652952663

Option 3 ID : 41652952662

Option 4 ID : 41652952665

Status : Not Answered

Chosen Option : --

Q.28

The coordination numbers of Co and Al in $[\text{Co}(\text{Cl})(\text{en})_2]\text{Cl}$ and $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$, respectively, are :
(en = ethane-1, 2-diamine)

- Options
1. 5 and 3
 2. 3 and 3
 3. 6 and 6
 4. 5 and 6

Question Type : MCQ
Question ID : 41652913464
Option 1 ID : 41652952637
Option 2 ID : 41652952635
Option 3 ID : 41652952634
Option 4 ID : 41652952636
Status : Answered
Chosen Option : 2

Q.29 The INCORRECT match in the following is :

- Options
1. $\Delta G^0 < 0, K > 1$
 2. $\Delta G^0 = 0, K = 1$
 3. $\Delta G^0 > 0, K < 1$
 4. $\Delta G^0 < 0, K < 1$

Question Type : MCQ
Question ID : 41652913469
Option 1 ID : 41652952654
Option 2 ID : 41652952657
Option 3 ID : 41652952656
Option 4 ID : 41652952655
Status : Answered
Chosen Option : 4

Q.30 Among the following, the INCORRECT statement about colloids is :

- Options
1. They can scatter light.
 2. They are larger than small molecules and have high molar mass.
 3. The osmotic pressure of a colloidal solution is of higher order than the true solution at the same concentration.
 4. The range of diameters of colloidal particles is between 1 and 1000 nm.

Question Type : MCQ
Question ID : 41652913475

Option 1 ID : **41652952679**
 Option 2 ID : **41652952681**
 Option 3 ID : **41652952680**
 Option 4 ID : **41652952678**
 Status : **Answered**
 Chosen Option : **2**

Section : Mathematics

Q.1

The derivative of $\tan^{-1}\left(\frac{\sin x - \cos x}{\sin x + \cos x}\right)$,

with respect to $\frac{x}{2}$, where $x \in \left(0, \frac{\pi}{2}\right)$ is:

- Options
1. 1
 2. $\frac{2}{3}$
 3. $\frac{1}{2}$
 4. 2

Question Type : **MCQ**
 Question ID : **41652913486**
 Option 1 ID : **41652952722**
 Option 2 ID : **41652952725**
 Option 3 ID : **41652952723**
 Option 4 ID : **41652952724**
 Status : **Answered**
 Chosen Option : **3**

Q.2

For an initial screening of an admission test, a candidate is given fifty problems to solve. If the probability that the candidate can solve any problem is $\frac{4}{5}$, then the probability that he is unable to solve less than two problems is:

- Options
1. $\frac{201}{5} \left(\frac{1}{5}\right)^{49}$
 2. $\frac{316}{25} \left(\frac{4}{5}\right)^{48}$
 3. $\frac{54}{5} \left(\frac{4}{5}\right)^{49}$
 4. $\frac{164}{25} \left(\frac{1}{5}\right)^{48}$

Question Type : **MCQ**
 Question ID : **41652913502**
 Option 1 ID : **41652952786**
 Option 2 ID : **41652952788**
 Option 3 ID : **41652952787**

Option 4 ID : **41652952789**
 Status : **Not Answered**
 Chosen Option : --

Q.3

A value of α such that

$$\int_{\alpha}^{\alpha+1} \frac{dx}{(x+\alpha)(x+\alpha+1)} = \log_e \left(\frac{9}{8} \right) \text{ is :}$$

- Options
1. -2
 2. $\frac{1}{2}$
 3. $-\frac{1}{2}$
 4. 2

Question Type : **MCQ**
 Question ID : **41652913490**
 Option 1 ID : **41652952740**
 Option 2 ID : **41652952738**
 Option 3 ID : **41652952741**
 Option 4 ID : **41652952739**
 Status : **Not Answered**
 Chosen Option : --

Q.4

Let $\alpha \in (0, \pi/2)$ be fixed. If the integral

$$\int \frac{\tan x + \tan \alpha}{\tan x - \tan \alpha} dx =$$

$A(x) \cos 2\alpha + B(x) \sin 2\alpha + C$, where C is a constant of integration, then the functions $A(x)$ and $B(x)$ are respectively :

- Options
1. $x + \alpha$ and $\log_e |\sin(x + \alpha)|$
 2. $x - \alpha$ and $\log_e |\sin(x - \alpha)|$
 3. $x - \alpha$ and $\log_e |\cos(x - \alpha)|$
 4. $x + \alpha$ and $\log_e |\sin(x - \alpha)|$

Question Type : **MCQ**
 Question ID : **41652913489**
 Option 1 ID : **41652952737**
 Option 2 ID : **41652952734**
 Option 3 ID : **41652952736**
 Option 4 ID : **41652952735**
 Status : **Not Answered**
 Chosen Option : --

Q.5

The angle of elevation of the top of a vertical tower standing on a horizontal plane is observed to be 45° from a point A on the plane. Let B be the point 30 m vertically above the point A. If the angle of elevation of the top of the tower from B be 30° , then the distance (in m) of the foot of the tower from the point A is :

Options

1. $15(3 + \sqrt{3})$
2. $15(5 - \sqrt{3})$
3. $15(3 - \sqrt{3})$
4. $15(1 + \sqrt{3})$

Question Type : MCQ

Question ID : 41652913504

Option 1 ID : 41652952795

Option 2 ID : 41652952797

Option 3 ID : 41652952794

Option 4 ID : 41652952796

Status : Marked For Review

Chosen Option : 2

Q.6 Let S be the set of all $\alpha \in \mathbb{R}$ such that the equation, $\cos 2x + \alpha \sin x = 2\alpha - 7$ has a solution. Then S is equal to :

Options

1. $\mathbb{R}$
2. $[1, 4]$
3. $[3, 7]$
4. $[2, 6]$

Question Type : MCQ

Question ID : 41652913503

Option 1 ID : 41652952790

Option 2 ID : 41652952792

Option 3 ID : 41652952793

Option 4 ID : 41652952791

Status : Not Answered

Chosen Option : --

Q.7 A plane which bisects the angle between the two given planes $2x - y + 2z - 4 = 0$ and $x + 2y + 2z - 2 = 0$, passes through the point :

Options

1. $(1, -4, 1)$
2. $(1, 4, -1)$
3. $(2, 4, 1)$

4. (2, -4, 1)

Question Type : MCQ

Question ID : 41652913498

Option 1 ID : 41652952770

Option 2 ID : 41652952773

Option 3 ID : 41652952771

Option 4 ID : 41652952772

Status : Not Attempted and Marked For Review

Chosen Option : --

Q.8

$$\lim_{x \rightarrow 0} \frac{x + 2\sin x}{\sqrt{x^2 + 2\sin x + 1} - \sqrt{\sin^2 x - x + 1}}$$

is :

Options

1. 6

2. 2

3. 3

4. 1

Question Type : MCQ

Question ID : 41652913485

Option 1 ID : 41652952721

Option 2 ID : 41652952719

Option 3 ID : 41652952720

Option 4 ID : 41652952718

Status : Not Answered

Chosen Option : --

Q.9

A group of students comprises of 5 boys and n girls. If the number of ways, in which a team of 3 students can randomly be selected from this group such that there is at least one boy and at least one girl in each team, is 1750, then n is equal to :

Options

1. 28

2. 27

3. 25

4. 24

Question Type : MCQ

Question ID : 41652913481

Option 1 ID : 41652952705

Option 2 ID : 41652952704

Option 3 ID : 41652952703

Option 4 ID : 41652952702

Status : Answered

Chosen Option : 3

Q.10

An ellipse, with foci at (0, 2) and (0, -2) and minor axis of length 4, passes through which of the following points ?

Options

1. $(\sqrt{2}, 2)$
2. $(2, \sqrt{2})$
3. $(2, 2\sqrt{2})$
4. $(1, 2\sqrt{2})$

Question Type : **MCQ**Question ID : **41652913497**Option 1 ID : **41652952768**Option 2 ID : **41652952766**Option 3 ID : **41652952769**Option 4 ID : **41652952767**Status : **Answered**Chosen Option : **1**

Q.11 The Boolean expression $\sim(p \Rightarrow (\sim q))$ is equivalent to :

Options

1. $p \wedge q$
2. $q \Rightarrow \sim p$
3. $p \vee q$
4. $(\sim p) \Rightarrow q$

Question Type : **MCQ**Question ID : **41652913505**Option 1 ID : **41652952800**Option 2 ID : **41652952799**Option 3 ID : **41652952801**Option 4 ID : **41652952798**Status : **Answered**Chosen Option : **3**

Q.12 A circle touching the x -axis at $(3, 0)$ and making an intercept of length 8 on the y -axis passes through the point :

Options

1. $(3, 10)$
2. $(3, 5)$
3. $(2, 3)$
4. $(1, 5)$

Question Type : **MCQ**Question ID : **41652913495**Option 1 ID : **41652952758**Option 2 ID : **41652952759**Option 3 ID : **41652952761**Option 4 ID : **41652952760**Status : **Not Answered**Chosen Option : **--****Q.13**

If ${}^{20}C_1 + (2^2) {}^{20}C_2 + (3^2) {}^{20}C_3 + \dots + (20^2) {}^{20}C_{20} = A(2^\beta)$, then the ordered pair (A, β) is equal to :

- Options
1. (420, 19)
 2. (420, 18)
 3. (380, 18)
 4. (380, 19)

Question Type : **MCQ**

Question ID : **41652913484**

Option 1 ID : **41652952717**

Option 2 ID : **41652952716**

Option 3 ID : **41652952714**

Option 4 ID : **41652952715**

Status : **Not Answered**

Chosen Option : --

Q.14 A value of $\theta \in (0, \pi/3)$, for which

$$\begin{vmatrix} 1+\cos^2\theta & \sin^2\theta & 4\cos 6\theta \\ \cos^2\theta & 1+\sin^2\theta & 4\cos 6\theta \\ \cos^2\theta & \sin^2\theta & 1+4\cos 6\theta \end{vmatrix} = 0, \text{ is :}$$

- Options
1. $\frac{\pi}{9}$
 2. $\frac{\pi}{18}$
 3. $\frac{7\pi}{24}$
 4. $\frac{7\pi}{36}$

Question Type : **MCQ**

Question ID : **41652913479**

Option 1 ID : **41652952695**

Option 2 ID : **41652952694**

Option 3 ID : **41652952696**

Option 4 ID : **41652952697**

Status : **Answered**

Chosen Option : **4**

Q.15 The equation of a common tangent to the curves, $y^2 = 16x$ and $xy = -4$, is :

- Options
1. $x - y + 4 = 0$
 2. $x + y + 4 = 0$
 3. $x - 2y + 16 = 0$
 4. $2x - y + 2 = 0$

Question Type : **MCQ**

Question ID : **41652913496**
 Option 1 ID : **41652952765**
 Option 2 ID : **41652952763**
 Option 3 ID : **41652952762**
 Option 4 ID : **41652952764**
 Status : **Answered**
 Chosen Option : **3**

Q.16 Let $z \in \mathbb{C}$ with $\text{Im}(z)=10$ and it satisfies

$$\frac{2z - n}{2z + n} = 2i - 1 \text{ for some natural number } n.$$

n. Then :

- Options**
1. $n = 20$ and $\text{Re}(z) = -10$
 2. $n = 40$ and $\text{Re}(z) = 10$
 3. $n = 40$ and $\text{Re}(z) = -10$
 4. $n = 20$ and $\text{Re}(z) = 10$

Question Type : **MCQ**
 Question ID : **41652913477**
 Option 1 ID : **41652952687**
 Option 2 ID : **41652952688**
 Option 3 ID : **41652952689**
 Option 4 ID : **41652952686**
 Status : **Not Answered**
 Chosen Option : --

Q.17 A triangle has a vertex at $(1, 2)$ and the mid points of the two sides through it are $(-1, 1)$ and $(2, 3)$. Then the centroid of this triangle is :

- Options**
1. $\left(1, \frac{7}{3}\right)$
 2. $\left(\frac{1}{3}, 2\right)$
 3. $\left(\frac{1}{3}, 1\right)$
 4. $\left(\frac{1}{3}, \frac{5}{3}\right)$

Question Type : **MCQ**
 Question ID : **41652913494**
 Option 1 ID : **41652952756**
 Option 2 ID : **41652952755**
 Option 3 ID : **41652952754**
 Option 4 ID : **41652952757**
 Status : **Not Answered**
 Chosen Option : --

Q.18

If $a_1, a_2, a_3, \dots$ are in A.P. such that $a_1 + a_7 + a_{16} = 40$, then the sum of the first 15 terms of this A.P. is :

- Options
1. 200
 2. 280
 3. 120
 4. 150

Question Type : **MCQ**

Question ID : **41652913483**

Option 1 ID : **41652952713**

Option 2 ID : **41652952712**

Option 3 ID : **41652952711**

Option 4 ID : **41652952710**

Status : **Answered**

Chosen Option : **3**

Q.19 If $[x]$ denotes the greatest integer $\leq x$, then the system of linear equations $[\sin\theta]x + [-\cos\theta]y = 0$
 $[\cot\theta]x + y = 0$

Options have infinitely many solutions if

1. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and has a unique

solution if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$.

has a unique solution if

2. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$.

has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$

3. and have infinitely many solutions if

$\theta \in \left(\pi, \frac{7\pi}{6}\right)$.

have infinitely many solutions if

4. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$.

Question Type : **MCQ**

Question ID : **41652913480**

Option 1 ID : **41652952701**

Option 2 ID : **41652952698**

Option 3 ID : **41652952700**

Option 4 ID : **41652952699**

Status : **Not Answered**

Chosen Option : **--**

Q.20 A straight line L at a distance of 4 units from the origin makes positive intercepts on the coordinate axes and the perpendicular from the origin to this line makes an angle of 60° with the line $x + y = 0$. Then an equation of the line L is :

- Options
1. $x + \sqrt{3}y = 8$
 2. $(\sqrt{3} + 1)x + (\sqrt{3} - 1)y = 8\sqrt{2}$
 3. $\sqrt{3}x + y = 8$
 4. $(\sqrt{3} - 1)x + (\sqrt{3} + 1)y = 8\sqrt{2}$

Question Type : **MCQ**

Question ID : **41652913493**

Option 1 ID : **41652952753**

Option 2 ID : **41652952750**

Option 3 ID : **41652952752**

Option 4 ID : **41652952751**

Status : **Not Answered**

Chosen Option : --

Q.21 Let $f(x) = 5 - |x - 2|$ and $g(x) = |x + 1|$, $x \in \mathbb{R}$. If $f(x)$ attains maximum value at α and $g(x)$ attains minimum value at β , then

$\lim_{x \rightarrow -\alpha\beta} \frac{(x - 1)(x^2 - 5x + 6)}{x^2 - 6x + 8}$ is equal to :

- Options
1. $1/2$
 2. $-3/2$
 3. $-1/2$
 4. $3/2$

Question Type : **MCQ**

Question ID : **41652913488**

Option 1 ID : **41652952730**

Option 2 ID : **41652952733**

Option 3 ID : **41652952731**

Option 4 ID : **41652952732**

Status : **Not Answered**

Chosen Option : --

Q.22 Let $\alpha \in \mathbb{R}$ and the three vectors

$$\vec{a} = \alpha \hat{i} + \hat{j} + 3\hat{k}, \quad \vec{b} = 2\hat{i} + \hat{j} - \alpha\hat{k}$$

and $\vec{c} = \alpha \hat{i} - 2\hat{j} + 3\hat{k}$. Then the set

$S = \{\alpha : \vec{a}, \vec{b} \text{ and } \vec{c} \text{ are coplanar}\}$

- Options
1. is singleton

2. is empty
3. contains exactly two positive numbers
4. contains exactly two numbers only one of which is positive

Question Type : **MCQ**
 Question ID : **41652913500**
 Option 1 ID : **41652952779**
 Option 2 ID : **41652952778**
 Option 3 ID : **41652952780**
 Option 4 ID : **41652952781**
 Status : **Answered**
 Chosen Option : **2**

Q.23 A person throws two fair dice. He wins Rs. 15 for throwing a doublet (same numbers on the two dice), wins Rs. 12 when the throw results in the sum of 9, and loses Rs. 6 for any other outcome on the throw. Then the expected gain/loss (in Rs.) of the person is :

- Options
1. $\frac{1}{2}$ gain
 2. $\frac{1}{4}$ loss
 3. $\frac{1}{2}$ loss
 4. 2 gain

Question Type : **MCQ**
 Question ID : **41652913501**
 Option 1 ID : **41652952784**
 Option 2 ID : **41652952785**
 Option 3 ID : **41652952783**
 Option 4 ID : **41652952782**
 Status : **Answered**
 Chosen Option : **3**

Q.24 The tangents to the curve $y = (x - 2)^2 - 1$ at its points of intersection with the line $x - y = 3$, intersect at the point :

- Options
1. $\left(\frac{5}{2}, 1\right)$
 2. $\left(-\frac{5}{2}, -1\right)$
 3. $\left(\frac{5}{2}, -1\right)$

4. $\left(-\frac{5}{2}, 1\right)$

Question Type : **MCQ**

Question ID : **41652913487**

Option 1 ID : **41652952728**

Option 2 ID : **41652952727**

Option 3 ID : **41652952726**

Option 4 ID : **41652952729**

Status : **Not Answered**

Chosen Option : --

Q.25 If α , β and γ are three consecutive terms of a non-constant G.P. such that the equations $\alpha x^2 + 2\beta x + \gamma = 0$ and $x^2 + x - 1 = 0$ have a common root, then $\alpha(\beta + \gamma)$ is equal to :

Options 1. 0

2. $\alpha\beta$

3. $\alpha\gamma$

4. $\beta\gamma$

Question Type : **MCQ**

Question ID : **41652913478**

Option 1 ID : **41652952693**

Option 2 ID : **41652952691**

Option 3 ID : **41652952692**

Option 4 ID : **41652952690**

Status : **Not Answered**

Chosen Option : --

Q.26 Let A, B and C be sets such that $\phi \neq A \cap B \subseteq C$. Then which of the following statements is not true ?

Options 1. $B \cap C \neq \phi$

2. If $(A - B) \subseteq C$, then $A \subseteq C$

3. $(C \cup A) \cap (C \cup B) = C$

4. If $(A - C) \subseteq B$, then $A \subseteq B$

Question Type : **MCQ**

Question ID : **41652913476**

Option 1 ID : **41652952682**

Option 2 ID : **41652952684**

Option 3 ID : **41652952683**

Option 4 ID : **41652952685**

Status : **Not Answered**

Chosen Option : --

Q.27 The general solution of the differential equation $(y^2 - x^3) dx - xy dy = 0$ ($x \neq 0$) is :
(where c is a constant of integration)

Options

1. $y^2 - 2x^2 + cx^3 = 0$
2. $y^2 + 2x^3 + cx^2 = 0$
3. $y^2 + 2x^2 + cx^3 = 0$
4. $y^2 - 2x^3 + cx^2 = 0$

Question Type : **MCQ**Question ID : **41652913492**Option 1 ID : **41652952749**Option 2 ID : **41652952748**Option 3 ID : **41652952746**Option 4 ID : **41652952747**Status : **Not Answered**

Chosen Option : --

Q.28 If the area (in sq. units) bounded by the parabola $y^2 = 4\lambda x$ and the line $y = \lambda x$, $\lambda > 0$, is $\frac{1}{9}$, then λ is equal to :

- Options
1. $2\sqrt{6}$
 2. 48
 3. 24
 4. $4\sqrt{3}$

Question Type : **MCQ**Question ID : **41652913491**Option 1 ID : **41652952743**Option 2 ID : **41652952745**Option 3 ID : **41652952744**Option 4 ID : **41652952742**Status : **Answered**

Chosen Option : 1

Q.29 The length of the perpendicular drawn from the point (2, 1, 4) to the plane containing the lines

$$\vec{r} = (\hat{i} + \hat{j}) + \lambda(\hat{i} + 2\hat{j} - \hat{k}) \quad \text{and}$$

$$\vec{r} = (\hat{i} + \hat{j}) + \mu(-\hat{i} + \hat{j} - 2\hat{k}) \quad \text{is:}$$

- Options
1. 3
 2. $\frac{1}{3}$
 3. $\sqrt{3}$
 4. $\frac{1}{\sqrt{3}}$

Question Type : **MCQ**Question ID : **41652913499**Option 1 ID : **41652952774**

Option 2 ID : **41652952776**
Option 3 ID : **41652952775**
Option 4 ID : **41652952777**
Status : **Not Answered**
Chosen Option : --

Q.30 The term independent of x in the expansion

of $\left(\frac{1}{60} - \frac{x^8}{81}\right) \cdot \left(2x^2 - \frac{3}{x^2}\right)^6$ is equal to :

- Options**
1. -72
 2. 36
 3. -36
 4. -108

Question Type : **MCQ**
Question ID : **41652913482**
Option 1 ID : **41652952707**
Option 2 ID : **41652952709**
Option 3 ID : **41652952708**
Option 4 ID : **41652952706**
Status : **Not Answered**
Chosen Option : --