

--	--	--	--	--

---

**Time : 3 Hours****MATHEMATICS****Subject Code**

H	4	7	5	4
---	---	---	---	---

**Total No. Of Questions : 36****(Printed Pages : 10)****Maximum Marks : 80**

---

- INSTRUCTIONS* :
- (i) The question paper consists of **36** questions.
  - (ii) All questions are compulsory.
  - (iii) Question numbers **1** to **8** are multiple choice type questions of *one* mark each.
  - (iv) Question numbers **9** to **16** are very short answer type questions of *one* mark each.
  - (v) Question numbers **17** to **22** are short answer type-I questions of *two* marks each.
  - (vi) Question numbers **23** to **28** are short answer type-II questions of *three* marks each.
  - (vii) Question numbers **29** to **34** are long answer type-I questions of *four* marks each.
  - (viii) Question numbers **35** to **36** are long answer type-II questions of *five* marks each.
  - (ix) There is no overall choice. However an internal choice has been provided in two questions of **4** marks each and 2 questions of **5** marks each.
  - (x) Use of calculator is not permitted.
  - (xi) Log tables will be supplied on request.
  - (xii) Graph should be drawn on the answer paper only.

1. The degree of the differential equation  $\left(\frac{d^2y}{dx^2}\right)^3 + xy = \left(\frac{dy}{dx}\right)^4$  is .....

- 1
- 2
- 3
- 4

2. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  are the two real functions defined by  $f(x) = 3x^2 + 1$  and  $g(x) = 1 - x$ , then  $(g \circ f)(-2)$  is .....

- 12
- 28
- -12
- -28

3. The value of  $[\hat{i} + \hat{j} \quad 2\hat{j} \quad 3\hat{k}]$  is .....

- 0
- 6
- 3
- 5

4. If  $\sin\left(\tan^{-1}(x) + \cot^{-1}\left(\frac{1}{3}\right)\right) = 1$ , then the value of  $x$  is .....

- 3
- -3
- $\frac{\pi}{2}$
- $\frac{1}{3}$

5. If  $R$  is a relation in the set  $A = \{3\}$  as  $R = \{(x, y) / x, y \in A \text{ and } x^2 + y + 4 \text{ is a perfect square}\}$ , then  $R$  is .....

- reflexive, symmetric but not transitive
- reflexive but neither symmetric nor transitive
- an equivalence relation
- symmetric but neither reflexive nor transitive

6. If  $\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} + X = \begin{bmatrix} 3 & 1 \\ 5 & 0 \end{bmatrix}$ , then the matrix  $X$  is .....

- $\begin{bmatrix} 6 & 5 \\ 4 & 4 \end{bmatrix}$
- $\begin{bmatrix} 0 & 5 \\ 4 & 2 \end{bmatrix}$
- $\begin{bmatrix} 0 & 5 \\ 4 & -2 \end{bmatrix}$
- $\begin{bmatrix} 0 & 5 \\ -4 & -2 \end{bmatrix}$

7. The value of  $\int_1^3 |x-3| dx$  is .....

- 1
- -2
- 2
- 0

8. The value of  $\int_{-1}^1 x^4 \sin^3 x dx$  is .....

- 1
- 0
- -1
- 2

9. Using determinants, show that the points (1, 3), (2, 2) and (0, 4) are collinear.

10. Find the slope of tangent to the curve  $2y = 3 - x^3$  at the point (1, 1).

11. Find the distance between the two planes :

$$x + y + 3z = 4 \text{ and } 2x + 2y + 6z = 10.$$

12. Find the area of parallelogram whose adjacent sides are given by the vectors

$$\bar{a} = \hat{i} + \hat{j} \text{ and } \bar{b} = 2\hat{i} + 3\hat{k}.$$

13. Find the principal value of  $\operatorname{cosec}^{-1}(-2)$ .

14. If

$$A = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}_{3 \times 1} \text{ and } B = [2 \ 0 \ 1]_{1 \times 3}$$

then find the matrix  $(AB)'$ , where  $(AB)'$  is the transpose of matrix  $(AB)$ .

15. The random variable X has the following probability distribution :

<b>X</b>	<b>P(X)</b>
0	K
1	2K
2	3K
3	4K

Find  $P(X < 2)$ .

16. If  $y = e^x + y^2$ , then find  $\frac{dy}{dx}$ .

17. Find the angle between the following pairs of lines :

$$\bar{r} = 2\hat{i} - 5\hat{j} + \hat{k} + \lambda(3\hat{i} + 2\hat{j} + 6\hat{k}) \text{ and}$$

$$\bar{r} = 7\hat{i} - 6\hat{k} + \mu(2\hat{i} + 2\hat{j} + \hat{k}).$$

18. If E and F are the events of a sample space S, such that  $P(F) \neq 0$ , then prove that :

$$P(E'/F) = 1 - P(E/F),$$

where E' is the complement of event E.

19. Form the differential equation of family of curves represented by  $A(y + A)^2 = x^3$ , by eliminating arbitrary constant A.
20. Let \* be a binary operation defined on set  $A = \{1, 2, 3, 6, 12\}$  as  $a * b = \text{H.C.F}\{a, b\}$ . Prepare composition table for the binary operation \*. Also, compute  $3 * (6 * 12)$ .

21. Prove that :

$$\tan^{-1} x - \tan^{-1} y = \tan^{-1} \left( \frac{x - y}{1 + xy} \right), \quad xy > -1.$$

22. If A(1, -2, 3) and B(-1, -4, 3) are the given points and  $\bar{d} = 3\hat{i} - 4\hat{j} + 5\hat{k}$  is a given vector, then find the scalar projection of vector  $\overline{AB}$  on  $\bar{d}$ .

23. Using properties of determinants, prove that :

$$\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} = (a+b+c)^3.$$

24. Find the general solution of the differential equation :

$$x^2 \frac{dy}{dx} = x^2 + xy - 2y^2.$$

25. One third of the students in a class are boys and the rest are girls. It is known that the probability of a girl getting first class is 0.32 and that a boy getting first class is 0.22. If a student chosen at random gets first class marks in the subject, what is the probability that the chosen student is a boy ?

26. If

$$y = (\sqrt{x})^{\sin 2x} + (\log 3x)^{\sqrt{x+1}},$$

find  $\frac{dy}{dx}$ .

27. Find the equation of the plane passing through the line of intersection of planes  $\vec{r} \cdot (2\hat{i} + \hat{j} - \hat{k}) = 3$  and  $\vec{r} \cdot (5\hat{i} - 3\hat{j} + 4\hat{k}) = -9$  and parallel to the line

$$\vec{r} = (\hat{i} + 3\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 4\hat{j} + 5\hat{k}).$$

28. Find :

$$\int \frac{x+5}{\sqrt{x^2+3x-7}} dx.$$

29. Prove that :

$$\int_0^{2a} f(x) dx = \int_0^a [f(x) + f(2a-x)] dx$$

Hence, show that :

$$\begin{aligned} \int_0^{2a} f(x) dx &= 2 \int_0^a f(x) dx, \quad \text{if } f(2a-x) = f(x) \\ &= 0, \quad \text{if } f(2a-x) = -f(x). \end{aligned}$$

30. Using integration, find the area of the smaller region bounded by the ellipse

$$\frac{x^2}{16} + \frac{y^2}{9} = 1 \quad \text{and the straight line } \frac{x}{4} + \frac{y}{3} = 1.$$

*Or*

Using integration, find the area of the smaller region enclosed between parabola  $y^2 = 16x$  and the line  $x - y + 3 = 0$ .

31. Solve the following Linear Programming Problem graphically :

$$\text{Minimize } Z = 5x + 7y$$

Subject to constraints :

$$2x + y \geq 8$$

$$x + 2y \geq 10$$

$$2x + 3y \leq 24$$

$$x \geq 0, y \geq 0.$$

32. If  $x = \sin t$ ,  $y = \sin(pt)$  prove that :

$$(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + p^2y = 0$$

where  $t$  is a parameter.

Or

If  $y = Ae^{-kx} \cos(px + 4)$ , prove that :

$$\frac{d^2y}{dx^2} + 2k\frac{dy}{dx} + (p^2 + k^2)y = 0.$$

33. If the function  $f(x)$  defined by :

$$\begin{aligned} f(x) &= \frac{1 + \sin x}{A \cos^2 x} & ; & \quad -\pi \leq x < \frac{-\pi}{2} \\ &= 3 \sin 2x + B & ; & \quad \frac{-\pi}{2} \leq x \leq 0 \\ &= \frac{e^{5x} - e^{3x}}{x} & ; & \quad 0 < x \leq \pi \end{aligned}$$

is continuous on  $[-\pi, \pi]$ , then find the values of A and B.

34. Find the inverse of matrix  $A = \begin{bmatrix} 1 & 2 & 5 \\ 1 & -1 & -1 \\ 2 & 3 & -1 \end{bmatrix}$ .

Hence, solve the system of equations :

$$x + 2y + 5z = 10$$

$$x - y - z = -2$$

$$2x + 3y - z = -11.$$

35. The perimeter of an isosceles triangle is 200 cm. If its base is changing at the rate 5 cm/sec, then find the rate at which the altitude is changing when the base is 40 cm.

*Or*

Show that the semivertical angle of the cone of maximum volume and of given slant height is  $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ .

36. Find :

$$\int \frac{2 \cos^2 x + \cos x}{(\sin x - 2)(\sin^2 x + 3)} dx.$$

*Or*

Find :

$$\int \cos^{-1}(\sqrt{x}) dx.$$