

CCE RF

ಕರ್ನಾಟಕ ಪ್ರೌಢ ಶಿಕ್ಷಣ ಪರೀಕ್ಷಾ ಮಂಡಳಿ, ಮಲ್ಲೇಶ್ವರಂ, ಬೆಂಗಳೂರು – 560 003

**KARNATAKA SECONDARY EDUCATION EXAMINATION BOARD, MALLESWARAM,
BANGALORE – 560 003**

ಎಸ್.ಎಸ್.ಎಲ್.ಸಿ. ಪರೀಕ್ಷೆ, ಮಾರ್ಚ್ / ಏಪ್ರಿಲ್ — 2015

S. S. L. C. EXAMINATION, MARCH/APRIL, 2015

ಮಾದರಿ ಉತ್ತರಗಳು

MODEL ANSWERS

ದಿನಾಂಕ : 06. 04. 2015]

ಸಂಕೇತ ಸಂಖ್ಯೆ : **81-E**

Date : 06. 04. 2015]

CODE No. : **81-E**

ವಿಷಯ : ಗಣಿತ

Subject : MATHEMATICS

(ಹೊಸ ಪಠ್ಯಕ್ರಮ / New Syllabus)

(ಶಾಲಾ ಅಭ್ಯರ್ಥಿ / Regular Fresh)

(ಇಂಗ್ಲಿಷ್ ಭಾಷಾಂತರ / English Version)

[ಪರಮಾವಧಿ ಅಂಕಗಳು : **80**[**Max. Marks : 80**

Qn. Nos.	Ans. Key	Value Points	Marks allotted
I. 1.	A	$\frac{8}{3}$	1
2.	C	1	1
3.	D	$\frac{1}{6}$	1
4.	B	2250	1
5.	B	- 2	1
6.	D	3	1
7.	A	5 units	1
8.	C	7	1

★ ★ ★

RF-126

★ ★ ★

[Turn over

Qn. Nos.	Value Points	Marks allotted	
II.			
9.	$210 = 2 \times 3 \times 5 \times 7$ $\begin{array}{r l} 2 & 210 \\ 3 & 105 \\ 5 & 35 \\ & 7 \end{array}$ $\frac{1}{2} + \frac{1}{2}$	1	
10.	$A \setminus B = \{2, 3, 4\}$	1	
11.	$C : V = \left(\frac{\sigma}{x}\right) \times 100$ $= \frac{4}{80} \times 100$ $\therefore CV = 5$	$\frac{1}{2}$ $\frac{1}{2}$	1
12.	$f(x) = x^2 - 4$ $f(4) = 4^2 - 4$ $= 16 - 4$ $\therefore f(4) = 12$	$\frac{1}{2}$ $\frac{1}{2}$	1
13.	$\frac{AX}{BX} = \frac{AY}{CY}$ $\frac{4}{BX} = \frac{1}{2}$ $\therefore BX = 8$	$\frac{1}{2}$ $\frac{1}{2}$	1
14.	$x = 90^\circ$	1	
III. 15.	<p>If possible let us assume $3 + \sqrt{5}$ is a rational number.</p> $3 + \sqrt{5} = \frac{p}{q}$ where $p, q \in \mathbb{Z}, q \neq 0$ $3 - \frac{p}{q} = -\sqrt{5}$ $\frac{3q - p}{q} = -\sqrt{5}$ $\Rightarrow -\sqrt{5}$ is a rational number. $\therefore \frac{3q - p}{q}$ is a rational number. But, $-\sqrt{5}$ is not a rational number \therefore Our supposition $3 + \sqrt{5}$ is a rational number is wrong. $\Rightarrow 3 + \sqrt{5}$ is an irrational number.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	2

Qn. Nos.	Value Points	Marks allotted
16.	<p>Number of students who have passed in Physics, $n(P) = 55$.</p> <p>Number of students who have passed in Mathematics, $n(M) = 67$.</p> <p>Number of students who have passed in both subjects, $n(P \cap M) = ?$</p> <p>Number of students in the classroom, $n(P \cup M) = 100$</p> $n(P) + n(M) = n(P \cup M) + n(P \cap M)$ $55 + 67 = 100 + n(P \cap M)$ $n(P \cap M) = 122 - 100$ $n(P \cap M) = 22$ <p>Number of students who have passed in Physics only</p> $= n(P) - n(P \cap M)$ $= 55 - 22$ $= 33.$ <p style="text-align: center;">OR</p> <p>$n(U) = 700$</p> <p>$n(A) = 200$</p> <p>$n(B) = 300$</p> <p>$n(A \cap B) = 100$</p> $n(A) + n(B) = n(A \cup B) + n(A \cap B)$ $200 + 300 = n(A \cup B) + 100$ $500 - 100 = n(A \cup B)$ <p>$n(A \cup B) = 400$</p> $n(A \cup B)^c = n(A^c \cap B^c)$ $= n[U \setminus (A \cup B)]$ $= n(U) - n(A \cup B)$ $= 700 - 400$ <p>$n(A \cup B)^c = 300$</p> <p>OR $n(A^c \cap B^c) = 300$</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2 $\frac{1}{2}$
17.	<p>Given digits : 1, 2, 3, 7, 8, 9</p> <p>a) 4-digit number can be formed in 6P_4 ways</p> ${}^6P_4 = 6 \times 5 \times 4 \times 3$ ${}^6P_4 = 360.$	$\frac{1}{2}$ $\frac{1}{2}$



Qn. Nos.	Value Points	Marks allotted								
18.	<p>b) Even numbers :</p> <table border="1" data-bbox="376 344 759 465"> <tr> <td><i>T</i></td> <td><i>H</i></td> <td><i>Ten</i></td> <td><i>U</i></td> </tr> <tr> <td>3P_1</td> <td>4P_1</td> <td>5P_1</td> <td>2P_1</td> </tr> </table> <p>Units place can be filled in 2P_1 ways Tens place can be filled in 5P_1 ways Hundreds place can be filled in 4P_1 ways Thousands place can be filled in 3P_1 ways Total number of ways = ${}^2P_1 \times {}^5P_1 \times {}^4P_1 \times {}^3P_1$ $= 2 \times 5 \times 4 \times 3$ $= 120.$</p> <p>$D = 35, n = ?$ $D = {}^nC_2 - n$ $35 = \frac{{}^nP_2}{2!} - n$ $35 = \frac{n(n-1)}{2} - n$ $35 = \frac{n^2 - n}{2} - n$ $35 = \frac{n^2 - 3n}{2}$ $n^2 - 3n - 70 = 0$ $n^2 - 10n + 7n - 70 = 0$ $n(n-10) + 7(n-10) = 0$ $n = 10$ OR $n = -7$ Neglecting $n = -7$ $n = 10$ \Rightarrow Number of sides = 10.</p>	<i>T</i>	<i>H</i>	<i>Ten</i>	<i>U</i>	3P_1	4P_1	5P_1	2P_1	$\frac{1}{2}$ $\frac{1}{2}$ 2 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2 $\frac{1}{2}$
<i>T</i>	<i>H</i>	<i>Ten</i>	<i>U</i>							
3P_1	4P_1	5P_1	2P_1							
19.	<p>$S = \{ (1, 1), (1, 2), \dots \dots (1, 6)$ $(6, 1), (6, 2), \dots \dots (6, 6) \}$ $\therefore n(S) = 36.$</p>	$\frac{1}{2}$								



RF-126



Qn. Nos.	Value Points	Marks allotted																		
	<p>a) Same number on both faces :</p> <p>Let A be the even.</p> $A = \{ (1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6) \}$ $n(A) = 6$ $P(A) = \frac{n(A)}{n(S)}$ $P(A) = \frac{6}{36}.$ <p>b) Both faces having multiples of five :</p> <p>Let B be the event.</p> $B = \{ (5, 5) \}$ $\therefore n(B) = 1$ $P(B) = \frac{n(B)}{n(S)}$ $\therefore P(B) = \frac{1}{36}.$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2																		
20.	<p>$X = 2, 4, 6, 8, 10$</p> <table border="1" data-bbox="375 1153 826 1451"> <thead> <tr> <th>X</th> <th>$D = X - \bar{x}$</th> <th>D^2</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>-4</td> <td>16</td> </tr> <tr> <td>4</td> <td>-2</td> <td>4</td> </tr> <tr> <td>6</td> <td>0</td> <td>0</td> </tr> <tr> <td>8</td> <td>2</td> <td>4</td> </tr> <tr> <td>10</td> <td>4</td> <td>16</td> </tr> </tbody> </table> $n = 5 \qquad \sum D^2 = 40$ $\bar{x} = \frac{\sum X}{n} = \frac{30}{5} = 6 \qquad \therefore \bar{x} = 6$ $\text{Variance} = \sigma^2 = \frac{\sum D^2}{n}$ $= \frac{40}{5}$ $\text{Variance} = \sigma^2 = 8.$	X	$D = X - \bar{x}$	D^2	2	-4	16	4	-2	4	6	0	0	8	2	4	10	4	16	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2
X	$D = X - \bar{x}$	D^2																		
2	-4	16																		
4	-2	4																		
6	0	0																		
8	2	4																		
10	4	16																		

Qn. Nos.	Value Points	Marks allotted												
24.	<p>By division algorithm for polynomials,</p> $P(x) = [g(x) \cdot q(x)] + r(x)$ $P(x) - r(x) = g(x) \cdot q(x)$ $P(x) + \{-r(x)\} = g(x) \cdot q(x)$ $x^2 + 2x - 3 \overline{) \begin{array}{r} x^4 + 2x^3 - 2x^2 + x - 1 \\ x^4 + 2x^3 - 3x^2 \\ \hline - x^2 + x - 1 \\ + 2x - 3 \\ \hline - x + 2 \end{array}}$ <p>$\therefore r(x) = -x + 2 \Rightarrow \{-r(x)\} = x - 2$</p> <p>Hence, we should add $(x - 2)$ to $P(x)$ so that the resulting polynomial is exactly divisible by $g(x)$.</p> $x^2 - 4x + 2 = 0$ $a = 1$ $b = -4$ $c = 2$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-4) \pm \sqrt{16 - 4(1)(2)}}{2(1)}$ $= \frac{4 \pm \sqrt{16 - 8}}{2}$ $= \frac{4 \pm \sqrt{8}}{2}$ $= \frac{4 \pm 2\sqrt{2}}{2}$ $= \frac{2(2 \pm \sqrt{2})}{2}$ $x = 2 \pm \sqrt{2}.$ <p>\therefore Roots are $2 + \sqrt{2}$ OR $2 - \sqrt{2}$.</p>	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; width: 50%; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td style="width: 50%;"></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td style="text-align: center; vertical-align: middle;">2</td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center; vertical-align: middle;">$\frac{1}{2}$</td> <td style="text-align: center; vertical-align: middle;">2</td> </tr> </table>	$\frac{1}{2}$		$\frac{1}{2}$		$\frac{1}{2}$		$\frac{1}{2}$	2	$\frac{1}{2}$		$\frac{1}{2}$	2
$\frac{1}{2}$														
$\frac{1}{2}$														
$\frac{1}{2}$														
$\frac{1}{2}$	2													
$\frac{1}{2}$														
$\frac{1}{2}$	2													


RF-126



[Turn over

Qn. Nos.	Value Points	Marks allotted								
25.	<div style="text-align: center;"> </div> <p>To prove : $PE = EQ$</p> <p>Proof : In ΔABD, $PE \parallel BD$ $\therefore PQ \parallel BC$</p> <p>$\therefore \frac{AP}{AB} = \frac{AQ}{AC} = \frac{PE}{BD}$ (i) (\because Thale's theorem)</p> <p>Similarly In ΔADC, $\frac{AE}{AD} = \frac{AQ}{AC} = \frac{EQ}{DC}$ (ii)</p> <p>From (i) and (ii) $\frac{PE}{BD} = \frac{EQ}{DC}$</p> <p>But, $BD = DC$ ($\because AD$ is the median)</p> <p>$\therefore PE = EQ$.</p>	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; width: 50px; text-align: center;">1/2</td> <td style="width: 50px;"></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1/2</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1/2</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1/2</td> <td style="text-align: right;">2</td> </tr> </table>	1/2		1/2		1/2		1/2	2
1/2										
1/2										
1/2										
1/2	2									
26.	<div style="text-align: center;"> </div> <p>In ΔABC, $\angle B = 90^\circ$</p> <p>$\therefore AC^2 = AB^2 + BC^2$</p> <p>$13^2 = 5^2 + BC^2$ (Pythagoras' theorem)</p> <p>$BC^2 = 169 - 25 = 144$</p> <p>$BC = 12$</p> <p>$\cos \theta = \frac{\text{Adjacent side}}{\text{Hypotenuse}} = \frac{BC}{AC}$</p> <p>$\therefore \cos \theta = \frac{12}{13}$</p> <p>$\tan \theta = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{AB}{BC}$</p> <p>$\therefore \tan \theta = \frac{5}{12}$.</p>	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; width: 50px; text-align: center;">1/2</td> <td style="width: 50px;"></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1/2</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1/2</td> <td style="text-align: right;">2</td> </tr> </table>	1/2		1/2		1/2	2		
1/2										
1/2										
1/2	2									




RF-126




Qn. Nos.	Value Points	Marks allotted
27.	<p>Let</p> $A = (x_1, y_1) = (3, 10)$ $B = (x_2, y_2) = (5, 2)$ $C = (x_3, y_3) = (14, 12)$ $AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(5 - 3)^2 + (2 - 10)^2}$ $AB = \sqrt{4 + 64} = \sqrt{68} \text{ units.}$ <p>Similarly</p> $BC = \sqrt{(14 - 5)^2 + (12 - 2)^2} = \sqrt{81 + 100} = \sqrt{181} \text{ units}$ $AC = \sqrt{(14 - 3)^2 + (12 - 10)^2} = \sqrt{121 + 4} = \sqrt{125} \text{ units}$ <p>Perimeter = $AB + BC + AC$</p> $P = \left(\sqrt{68} + \sqrt{181} + \sqrt{125} \right) \text{ units.}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2
28.	<p>$r = 3 \text{ cm}, d = 8 \text{ cm.}$</p>  <p>PA and PB are tangents.</p> <p>Drawing circle of radius 3 cm</p> <p>Bisecting OP</p> <p>Drawing tangents PA, PB.</p>	$\frac{1}{2}$ $\frac{1}{2}$ 1 2



Qn. Nos.	Value Points	Marks allotted
29.	<p>Scale : 20 m = 1 cm 40 m = 2 cm 60 m = 3 cm 80 m = 4 cm 100 m = 5 cm 150 m = 7.5 cm</p> <p style="text-align: right;">Calculation Field drawing</p> 	$\frac{1}{2}$ $1\frac{1}{2}$ 2
30.	<p>$F = 6$ $V = 8$ $E = 12$</p> $F + V = E + 2$ $6 + 8 = 12 + 2$ $14 = 14.$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 2
IV. 31.	<p>$T_3 = \frac{1}{7}, T_7 = \frac{1}{5}$</p> <p>$\therefore T_3$ and T_7 of A.P. are 7, 5</p> $d = \frac{T_p - T_q}{p - q}$ $= \frac{T_7 - T_3}{7 - 3} = \frac{5 - 7}{4} = \frac{-2}{4} = -\frac{1}{2}$ <p>$\therefore d = -\frac{1}{2}$</p> <p>$T_3 = 7$</p> <p>$a + 2d = 7$</p>	1




Qn. Nos.	Value Points	Marks allotted
33.	<p>Let the number of years be x</p> <p>\therefore Kavya's and Karthik's ages are $(11 + x)$ and $(14 + x)$</p> <p>Product of their ages = 304</p> $(11 + x)(14 + x) = 304$ $154 + 11x + 14x + x^2 - 304 = 0$ $x^2 + 25x - 150 = 0$ $x^2 + 30x - 5x - 150 = 0$ $x(x + 30) - 5(x + 30) = 0$ $x = 5 \quad \text{OR} \quad -30$ <p>Neglecting $x = -30$</p> $x = 5.$ <p>i.e. After 5 years, product of their ages will be 304.</p> <div style="text-align: center;">  </div> <p>In ΔABC,</p> $\hat{A} = 90^\circ, \quad AB = x, \quad \hat{B} = 45^\circ \quad \Rightarrow \quad \hat{C} = 45^\circ$ $\Rightarrow \quad AB = AC = x$ $BD = CD = \frac{BC}{2}$ $BC^2 = AC^2 + AB^2 \quad \dots \text{Pythagoras' theorem}$ $= x^2 + x^2$ $BC^2 = 2x^2$ $\therefore \quad BC = x\sqrt{2}$ $AD^2 = CD \cdot BD$ $= \frac{1}{2} BC \cdot \frac{1}{2} BC$ $= \left(\frac{BC}{2}\right)^2$	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>3</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>



RF-126




Qn. Nos.	Value Points	Marks allotted
	$= \left(\frac{x\sqrt{2}}{2} \right)^2$ $= \frac{x^2 \cdot 2}{4}$ $\therefore AD^2 = \frac{x^2}{2}$ $\therefore AD = \frac{x}{\sqrt{2}}$	1 1/2 3
	<p style="text-align: center;">OR</p>  <p>To prove : $\frac{1}{p^2} = \frac{1}{a^2} + \frac{1}{b^2}$</p> <p>Proof : In ΔABC $CD^2 = AD \cdot BD$ $p^2 = AD \cdot BD$ (i)</p> <p>$CB^2 = AB \cdot AD$ $a^2 = AB \cdot AD$ $\therefore \frac{1}{a^2} = \frac{1}{AB \cdot BD}$ (ii)</p> <p>$AC^2 = AB \cdot AD$ $b^2 = AB \cdot AD$ $\therefore \frac{1}{b^2} = \frac{1}{AB \cdot AD}$ (iii)</p> <p>Adding (ii) and (iii) $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{AB \cdot BD} + \frac{1}{AB \cdot AD}$ $= \frac{1}{AB} \left(\frac{1}{BD} + \frac{1}{AD} \right)$ $= \frac{1}{AB} \left(\frac{AD + BD}{BD \cdot AD} \right) = \frac{1}{AB} \cdot \frac{AB}{p^2}$ [from (i)]</p> <p>$\therefore \frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2}$</p>	1/2 1/2 1/2 1 3

★ ★ ★

RF-126

★ ★ ★

[Turn over

Qn. Nos.	Value Points	Marks allotted
35.	<div style="text-align: center;">  </div> <p><i>Data :</i> O is the centre of the circle B is the external point BP and BQ are the tangents.</p> <p><i>To prove :</i> $BP = BQ$</p> <p><i>Proof :</i> In $\triangle BOP$, $\hat{BPO} = 90^\circ$ Similarly in $\triangle BOQ$, $\hat{BQO} = 90^\circ$ (\because radius, tangents at the point of contact are perpendicular)</p> <p>In 2 right angled triangles BOP and BOQ Hypotenuse $BO =$ Hypotenuse BO Common side $OP = OQ$... radii $\therefore \triangle BOP \cong \triangle BOQ$ $\therefore BP = BQ$.</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3
36.	<p>CSA of cylinder = $\frac{1}{3}$ TSA of cylinder</p> $2\pi rh = \frac{1}{3} \cdot 2\pi r (r + h)$ $= \frac{1}{3} \times 462$ $2\pi rh = 154$ <p>TSA of cylinder = $2\pi r (r + h)$</p> $462 = 2\pi r^2 + 2\pi rh$ $462 = 2\pi r^2 + 154$ $462 - 154 = 2\pi r^2$ $2\pi r^2 = 308$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

★ ★ ★

RF-126

★ ★ ★

[Turn over


Qn. Nos.	Value Points	Marks allotted
	$r^2 = \frac{308}{2 \times \frac{22}{7}}$ $= \frac{308 \times 7}{44}$ $r^2 = 49$ $r = 7 \text{ cm.}$ <p style="text-align: center;">OR</p> $r_1 = 12 \text{ cm, } h_1 = 20 \text{ cm, } r_2 = 3 \text{ cm, } h_2 = ?$ <p>We know,</p> $\frac{r_1}{r_2} = \frac{h_1}{h_2}$ $\frac{12}{3} = \frac{20}{h_2}$ $12h_2 = 60 \quad \therefore h_2 = 5 \text{ cm}$ <p>Volume of frustum = $\frac{1}{3} \pi h (r_1^2 + r_2^2 + r_1 r_2)$</p> $= \frac{1}{3} \times \frac{22}{7} \times 15 (12^2 + 3^2 + 12 \times 3)$ $= \frac{110}{7} (144 + 9 + 36)$ $= \frac{110}{7} \times 189$ <p>Volume of frustum = 2970 cubic cm.</p>	$\frac{1}{2}$ $\frac{1}{2}$ 3 $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 3
V. 37.	<p>Let the three consecutive terms of the A.P. be $(a - d), a, (a + d)$</p> <p>Sum = 6</p> $a - d + a + a + d = 6$ $3a = 6$ $\therefore a = 2$ <p>Product = -120</p> $(a - d) \cdot a (a + d) = -120$ $(a^2 - d^2) a = -120$ $(2^2 - d^2) 2 = -120$ $4 - d^2 = -60$ $-d^2 = -64$	$\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$



RF-126



Qn. Nos.	Value Points	Marks allotted
	$d^2 = 64 \quad \therefore d = \pm 8$ If $a = 2, d = 8$ then the three numbers are $-6, 2, 10$ If $a = 2, d = -8$, then the three numbers are $10, 2, -6$	$\frac{1}{2}$ 1 4
	OR Let the three consecutive terms be $\frac{a}{r}, a, ar$ Product = 216 $\frac{a}{r} \cdot a \cdot ar = 216$ $a^3 = 216 \quad \therefore a = 6$	1
	Sum of their products in pairs = 156 $\left(\frac{a}{r} \cdot a\right) + (a \cdot ar) + \left(\frac{a}{r} \cdot ar\right) = 156$	$\frac{1}{2}$
	$\frac{a^2}{r} + a^2r + a^2 = 156$ $\frac{36}{r} + 36r + 36 = 156$	$\frac{1}{2}$
	$\frac{36}{r} + 36r = 120$ $\frac{36 + 36r^2}{r} = 120$	
	$36r^2 - 120r + 36 = 0$ $3r^2 - 10r + 3 = 0$	
	$r = 3$ or $\frac{1}{3}$	$\frac{1}{2} + \frac{1}{2}$
	a) If $a = 6, r = 3$, the 3 consecutive numbers are 2, 6, 18 b) If $a = 6, r = \frac{1}{3}$, the 3 consecutive numbers are 18, 6, 2	$\frac{1}{2}$ $\frac{1}{2}$ 4

Qn. Nos.	Value Points	Marks allotted
38.	<div style="text-align: center;">  </div> <p><i>Data :</i> In $\triangle ABC$, $\hat{B} = 90^\circ$</p> <p><i>To prove :</i> $AC^2 = AB^2 + BC^2$</p> <p><i>Constn. :</i> Draw $BD \perp AC$</p> <p><i>Proof :</i> In 2 triangles ABC, ADB $\hat{A}BC = 90^\circ$, $\hat{ADB} = 90^\circ$ \hat{BAD} ... common angle $\therefore \triangle ABC \sim \triangle ADB$ $\Rightarrow \frac{AB}{AD} = \frac{AC}{AB}$ $AB^2 = AC \cdot AD$ (i)</p> <p>In 2 triangles ABC and BDC $\hat{ABC} = \hat{BDC} = 90^\circ$ \hat{ACB} is common $\therefore \triangle ABC \sim \triangle BDC$ $\Rightarrow \frac{BC}{DC} = \frac{AC}{BC}$ $\therefore BC^2 = AC \cdot DC$ (ii)</p> <p>Adding (i) and (ii) $AB^2 + BC^2 = (AC \cdot AD) + (AC \cdot DC)$ $= AC (AD + DC)$ $AB^2 + BC^2 = AC^2$ OR $AC^2 = AB^2 + BC^2$</p>	<div style="text-align: right;"> $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 </div>

★ ★ ★

RF-126

★ ★ ★


Qn. Nos.	Value Points	Marks allotted								
39.	<p>$d = 9 \text{ cm}, R = 4 \text{ cm}, r = 2 \text{ cm}, R - r = 2 \text{ cm}.$</p> <p>Drawing AB and marking mid-point M</p> <p>Drawing circles C_1, C_2 and C_3</p> <p>Joining BK, BL, PQ, RS</p> <p>Measuring and writing the length of tangents = 8.7 cm</p> <div style="text-align: center;"> </div> <p>Required tangents $PQ = RS = 8.7 \text{ cm}.$</p>	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-right: 1px solid black; text-align: center;">1</td> <td style="width: 50%;"></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">$1\frac{1}{2}$</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">1</td> <td></td> </tr> <tr> <td style="border-right: 1px solid black; text-align: center;">$\frac{1}{2}$</td> <td style="text-align: center; vertical-align: middle;">4</td> </tr> </table>	1		$1\frac{1}{2}$		1		$\frac{1}{2}$	4
1										
$1\frac{1}{2}$										
1										
$\frac{1}{2}$	4									



RF-126



[Turn over

Qn. Nos.	Value Points	Marks allotted																											
40.	$x^2 + x - 6 = 0$ Let $y = x^2 + x - 6$ <table border="1" style="margin: 10px auto;"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>-1</td> <td>2</td> <td>-2</td> <td>3</td> <td>-3</td> <td>-4</td> </tr> <tr> <td>y</td> <td>-6</td> <td>-4</td> <td>-6</td> <td>0</td> <td>-4</td> <td>6</td> <td>0</td> <td>6</td> </tr> </table>  <p style="text-align: right;">Table Drawing parabola Identifying roots</p> <p style="text-align: right;">Roots are 2 or -3</p>	x	0	1	-1	2	-2	3	-3	-4	y	-6	-4	-6	0	-4	6	0	6	<table style="border-collapse: collapse;"> <tr> <td style="border-right: 1px solid black; padding: 5px;">2</td> <td style="padding: 5px;"> </td> <td style="padding: 5px;">4</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 5px;">1</td> <td style="padding: 5px;"> </td> <td style="padding: 5px;"></td> </tr> <tr> <td style="border-right: 1px solid black; padding: 5px;">1</td> <td style="padding: 5px;"> </td> <td style="padding: 5px;"></td> </tr> </table>	2		4	1			1		
x	0	1	-1	2	-2	3	-3	-4																					
y	-6	-4	-6	0	-4	6	0	6																					
2		4																											
1																													
1																													
	<p><i>Alternate Method :</i></p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Table</td> <td style="text-align: right;">1 + 1</td> </tr> <tr> <td>Parabola</td> <td style="text-align: right;">1/2</td> </tr> <tr> <td>Straight line</td> <td style="text-align: right;">1/2</td> </tr> <tr> <td>Drawing perpendiculars and identifying roots</td> <td style="text-align: right;">1/2 + 1/2</td> </tr> </table>	Table	1 + 1	Parabola	1/2	Straight line	1/2	Drawing perpendiculars and identifying roots	1/2 + 1/2	4																			
Table	1 + 1																												
Parabola	1/2																												
Straight line	1/2																												
Drawing perpendiculars and identifying roots	1/2 + 1/2																												

